


Cell Metabolism

Intercellular Mitochondria Transfer to Macrophages Regulates White Adipose Tissue Homeostasis and Is Impaired in Obesity

Graphical Abstract



Authors

Jonathan R. Brestoff, Craig B. Wilen, John R. Moley, ..., Herbert W. Virgin, Andrew E. Gelman, Steven L. Teitelbaum

Correspondence

teitelbs@wustl.edu

In Brief

Brestoff et al. show that adipose-tissue-resident macrophages acquire mitochondria from neighboring adipocytes in a heparan-sulfate-dependent process that is impaired in obesity. Genetic disruption of mitochondria uptake by macrophages reduces energy expenditure and exacerbates diet-induced obesity in mice, indicating that intercellular mitochondria transfer to macrophages mediates systemic metabolic homeostasis.

Highlights

- Adipocytes transfer their mitochondria to macrophages *in vivo*
- Mitochondria transfer from adipocytes to macrophages is decreased in obesity
- Mitochondria uptake by macrophages is mediated by heparan sulfates
- Mice that lack heparan sulfates on macrophages exhibit metabolic dysfunction



Article

Intercellular Mitochondria Transfer to Macrophages Regulates White Adipose Tissue Homeostasis and Is Impaired in Obesity

Jonathan R. Brestoff,¹ Craig B. Wilen,² John R. Moley,¹ Yongjia Li,¹ Wei Zou,¹ Nicole P. Malvin,¹ Marina N. Rowen,¹ Brian T. Saunders,¹ Hongming Ma,³ Madison R. Mack,^{3,4,5,6} Barry L. Hykes, Jr.,¹ Dale R. Balce,^{1,7} Anthony Orvedahl,⁸ Jesse W. Williams,¹ Nidhi Rohatgi,¹ Xiaoyan Wang,¹ Michael R. McAllaster,¹ Scott A. Handley,¹ Brian S. Kim,^{1,3,4,6} John G. Doench,⁹ Bernd H. Zinselmeyer,¹ Michael S. Diamond,^{1,3,10} Herbert W. Virgin,^{1,7,11} Andrew E. Gelman,^{1,12} and Steven L. Teitelbaum^{1,3,13,*}

¹Department of Pathology and Immunology, Washington University School of Medicine, St. Louis, MO 63110, USA

²Department of Laboratory Medicine and Department of Immunobiology, Yale University School of Medicine, New Haven, CT, USA

³Department of Medicine, Washington University School of Medicine, St. Louis, MO 63110, USA

⁴Center for the Study of Itch, Washington University School of Medicine, St. Louis, MO 63110, USA

⁵Division of Biology and Biomedical Sciences, Washington University School of Medicine, St. Louis, MO 63110, USA

⁶Department of Anesthesiology, Washington University School of Medicine, St. Louis, MO 63110, USA

⁷Vir Biotechnology, San Francisco, CA 94158, USA

⁸Department of Pediatrics, Washington University School of Medicine, St. Louis, MO 63110, USA

⁹Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA

¹⁰Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, MO 63110, USA

¹¹Department of Medicine, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA

¹²Department of Surgery, Washington University School of Medicine, St. Louis, MO 63110, USA

¹³Lead Contact

*Correspondence: teitelbs@wustl.edu

<https://doi.org/10.1016/j.cmet.2020.11.008>

SUMMARY

Recent studies suggest that mitochondria can be transferred between cells to support the survival of metabolically compromised cells. However, whether intercellular mitochondria transfer occurs in white adipose tissue (WAT) or regulates metabolic homeostasis *in vivo* remains unknown. We found that macrophages acquire mitochondria from neighboring adipocytes *in vivo* and that this process defines a transcriptionally distinct macrophage subpopulation. A genome-wide CRISPR-Cas9 knockout screen revealed that mitochondria uptake depends on heparan sulfates (HS). High-fat diet (HFD)-induced obese mice exhibit lower HS levels on WAT macrophages and decreased intercellular mitochondria transfer from adipocytes to macrophages. Deletion of the HS biosynthetic gene *Ext1* in myeloid cells decreases mitochondria uptake by WAT macrophages, increases WAT mass, lowers energy expenditure, and exacerbates HFD-induced obesity *in vivo*. Collectively, this study suggests that adipocytes and macrophages employ intercellular mitochondria transfer as a mechanism of immunometabolic crosstalk that regulates metabolic homeostasis and is impaired in obesity.

INTRODUCTION

Obesity is an increasingly prevalent metabolic disease that affects 38% of adults and 16% of children and adolescents and is an independent risk factor for the development of many other disorders such as type 2 diabetes, cardiovascular diseases, and cancer (Flegal et al., 2016; Ogden et al., 2016; Patterson et al., 2004). Although obesity pathogenesis is driven by chronic positive energy balance, where caloric intake exceeds caloric expenditure, recent studies have revealed that the immune system modulates neuroendocrine pathways that govern not only food intake but also energy expenditure (Brestoff and Artis, 2015; Lee et al., 2018; Reilly and Saltiel, 2017; Sun et al., 2011; Wellen and Hotamisligil, 2003). In particular,

macrophages in white adipose tissue (WAT) regulate metabolic homeostasis by promoting glucose utilization by adipocytes, regulating lipid storage and release, and increasing energy expenditure through modulation of sympathetic tone to WAT (Camell et al., 2017; Flaherty et al., 2019; Pirzgalska et al., 2017; Qiu et al., 2014; Rao et al., 2014; Wu et al., 2011; Wynn et al., 2013). In the context of obesity, various cytokines activate macrophages in WAT to promote a chronic inflammation that dysregulates glucose, lipid, and energy metabolism (Hotamisligil, 2017; Larabee et al., 2020; McLaughlin et al., 2017; Reilly and Saltiel, 2017). Despite these critical advances, we lack a complete understanding of how macrophages and adipocytes interact with each other to maintain metabolic homeostasis.

In 2006, it was reported that mtDNA-deficient ρ^0 cells, which lack functionally competent mitochondria, can capture purified mitochondria from supernatants and that this process promotes cell proliferation (Spees et al., 2006). This phenomenon of mitochondria uptake has been confirmed in various ρ^0 cell lines and can restore normal mitochondrial respiration in ρ^0 cells (Jackson and Krasnodembskaya, 2017; Kim et al., 2018; Sinha et al., 2016). It is known that cells can release functional mitochondria that are then captured by recipient cells (Jiang et al., 2016; Kitani et al., 2014; Maeda and Fadeel, 2014; Scozzi et al., 2019; Torralba et al., 2016). It is also reported that cells can release extracellular vesicles (EVs) that contain mitochondria, with recipient cells acquiring mitochondria through an EV-cell fusion event (Boudreau et al., 2014; Torralba et al., 2016). In both mice and humans, mitochondria and EV-associated mitochondria circulate in blood (Al Amir Dache et al., 2020; Boudreau et al., 2014; Chiu et al., 2003; Pollara et al., 2018; Scozzi et al., 2019). Intercellular mitochondria transfer has been implicated in several disease processes, including ischemic stroke (Hayakawa et al., 2016), inflammatory pain (Raoof et al., 2020), cancer cell growth (Chang et al., 2019; Dong et al., 2017; Griessinger et al., 2017; Rebbeck et al., 2011), acute lung injury repair (Islam et al., 2012), hypoxia-induced pulmonary hypertension (Zhu et al., 2016), and allograft rejection (Lin et al., 2000; Pollara et al., 2018; Scozzi et al., 2019). However, it remains unknown whether intercellular mitochondria transfer occurs in adipose tissues or whether this process regulates systemic metabolic homeostasis.

Here, we employed an adipocyte-specific mitochondria reporter (MitoFat) mouse and demonstrate that macrophages in WAT acquire mitochondria from adipocytes *in vivo*. This adipocyte-to-macrophage mitochondrial transfer axis defines a distinct macrophage subpopulation in WAT and is severely decreased in high-fat diet (HFD) obesity due to reduced mitochondria uptake by macrophages. A genome-wide CRISPR-Cas9 screen revealed that mitochondria uptake depends on the heparan sulfate (HS) biosynthetic pathway, including the genes *Exostosin 1* (*Ext1*) and its heterodimer *Ext2*, which have been linked to the maintenance of glucose and lipid homeostasis in mice and humans (Busse et al., 2007; Liu et al., 2013; Mooij et al., 2015; Sladek et al., 2007). Epididymal (e)WAT macrophage HS levels are decreased in obese mice, and IFN- γ plus LPS treatment downregulates expression of HS synthetic enzymes and impairs mitochondria uptake despite promoting bead phagocytosis. Myeloid-cell-specific deletion of *Ext1*, which is required for HS synthesis (Busse et al., 2007; Lin et al., 2000; Sasisekharan and Venkataraman, 2000), impairs mitochondria uptake by macrophages in WAT and leads to dysregulated energy homeostasis characterized by lower energy expenditure, accumulation of WAT mass, glucose intolerance, and increased susceptibility to diet-induced obesity. Collectively, these observations suggest that intercellular mitochondria transfer represents an unappreciated mechanism of cellular communication that may regulate systemic metabolic homeostasis.

RESULTS

Macrophages in WAT Acquire Mitochondria from Other Cell Types *In Vivo*

To test whether intercellular mitochondria transfer occurs *in vivo*, we performed bone marrow transplants of congenic

CD45.1 wild-type (WT) bone marrow into host CD45.2 mitochondria reporter mice that express the fluorescent protein Dendra2 attached to the mitochondria targeting sequence of cytochrome c, complex VIII (mtD2) (Pham et al., 2012) (Figure 1A). As a control, WT CD45.1 bone marrow was transplanted into WT CD45.2 recipients. After 12 weeks of engraftment, the stromal vascular fraction (SVF) was isolated from eWAT, inguinal (i)WAT, and brown adipose tissue (BAT) for flow cytometric analysis. The vast majority of the immune cells in these organs were WT donor-derived CD45.1⁺ CD45.2⁻ cells, with chimerism of 77% in eWAT, 88% in iWAT, and 72% in BAT (Figure S1A). We then gated on WT CD45.2⁺ CD45.1⁻ donor-derived immune cells in eWAT and found that nearly 40% were mtD2⁺ when isolated from mtD2 hosts (Figure 1B). The majority of these donor-derived mtD2⁺ cells were macrophages (65.5%), with other myeloid cells such as monocytes, CD11c⁺ cells, eosinophils, and neutrophils representing most of the remainder (Figure 1C; gating strategy shown in Figure S1B). Next, we gated on WT CD45.1⁺ CD45.2⁻ F4/80⁺ CD64⁺ donor-derived macrophages and found that approximately 60% were mtD2⁺ when isolated from mtD2 hosts compared with approximately 3% from WT hosts ($p < 0.001$; Figure 1D). The proportion of macrophages that were mtD2⁺ was significantly higher than that of monocytes, CD11c⁺ cells, eosinophils, neutrophils, CD4⁺ cells, or other undefined cell types (Figure 1E). Further, the proportion of macrophages that are mtD2⁺ was significantly higher in eWAT and iWAT than in BAT (Figure 1F). Collectively, these bone marrow transplant data suggest that macrophages in WAT and, to a lesser degree BAT, acquire mitochondria from host cells.

A limitation of flow cytometry is that it cannot distinguish between surface binding and internalization of host-cell-derived mitochondria. To determine if mitochondria are internalized *in vivo*, we isolated macrophages from eWAT of WT mice, labeled them with CytoTracker Orange (CMTMR), adoptively transferred these cells into mtD2 host mice, and then performed intravital 2-photon microscopy on host eWAT (Figure 2A). This imaging technique allowed us to capture images approximately 25–30 μ m deep within eWAT of live mice. Host eWAT revealed large adipocytes (dashed circle), host-derived mtD2⁺ mitochondria, and CMTMR⁺ donor macrophages adjacent to host adipocytes (Figure 2B). This indicates that adoptively transferred eWAT macrophages can relocate to their tissue of origin, as has been observed with eosinophils (Wu et al., 2011) and group 2 innate lymphoid cells (Brestoff et al., 2015). Critically, CMTMR⁺ donor macrophages contained internalized host-derived mitochondria and appeared to exhibit interactions with host-derived mitochondria at the plasma membrane (Figure 2C). Three-dimensional reconstructions with positional mapping confirmed that host-cell-derived mitochondria can be found within CMTMR⁺ donor macrophages (Figure 2D). In addition, time-lapse videography revealed a mitochondria uptake event, trafficking of host-cell-derived mitochondria within CMTMR⁺ donor macrophages, and apparent fusion and fission involving internalized mitochondria (Figure 2E; Video S1). These data confirm that macrophages in WAT can internalize mitochondria from other cell types *in vivo*.




Figure 1. Macrophages Acquire Mitochondria from Other Cell Types In Vivo

(A) Experimental design.

(B) Frequency of live-donor-derived WT immune cells that are mtD2⁺ in recipient epididymal white adipose tissue (eWAT). Pre-gated on live CD45.1⁺ CD45.2⁻ cells.

(C) Cell types represented within the population of WT donor cells that are mtD2⁺.

(D) Frequencies of WT donor-derived macrophages that are mtD2⁺ in eWAT. Pre-gated on live CD45.1⁺ CD45.2⁻ CD64⁺ F4/80⁺ macrophages in eWAT.

(E) Frequencies of the indicated donor-derived immune cells that are mtD2⁺ in eWAT.

(F) Frequencies of donor-derived macrophages that are mtD2⁺ in eWAT compared with inguinal (i)WAT and brown adipose tissue (BAT).

Data represent 2–4 independent experiments with n = 4–8 mice/group combined and are expressed as mean ± SEM. Student's t tests (B and D), one-way ANOVA with Fisher's post-hoc test compared with macrophages (E), and two-way ANOVA with Fisher's post-hoc test (F); *p < 0.05, **p < 0.01, ***p < 0.001. See also Figure S1.

Adipocytes Transfer Mitochondria to Macrophages In Vivo, Defining a Distinct Macrophage Subpopulation

Given the abundance of adipocytes in WAT, we hypothesized that adipocytes might transfer their mitochondria to macrophages. To test this, we generated adipocyte-specific mitochondria reporter mice (MitoFat) by crossing *mtD2^{Flox/Flox}* (Pham et al., 2012) to *Adipoq^{Cre/+}* mice. We found that approximately 40% of macrophages in gonadal (g)WAT contained mitochondria that were derived from adipocytes (Figures 3A and 3B). To confirm that adipocytes transfer their mitochondria to macrophages, we adoptively transferred CMTMR-labeled eWAT macrophages into MitoFat mice and performed intravital 2-photon microscopy of host eWAT. Three-dimensional reconstructions with positional mapping confirmed that adipocyte-derived mitochondria can be identified within CMTMR⁺ donor macrophages (Figure 3C). These observations define an adipocyte-to-macrophage mitochondrial transfer axis that occurs in WAT *in vivo*.

To test whether intercellular mitochondria transfer from adipocytes to macrophages is associated with alterations in the phenotype of WAT macrophages, we isolated WAT SVF from MitoFat mice and directly compared macrophages that had (mtD2⁺) or had not (mtD2⁻) received mitochondria from adipocytes (Figure 3D). First, we found that mtD2⁺ macrophages had modestly (15%) but significantly higher mitochondrial mass, as indicated by the charge-independent MitoID-Red mitochondria dye, than mtD2⁻ macrophages from the same samples (Figure 3E). Overall mitochondrial membrane potential, as indicated by the charge-dependent CMX Rosamine dye, was

decreased in mtD2⁺ versus mtD2⁻ macrophages (Figure 3F). MitoSOX staining, an indicator of mitochondria reactive oxygen species including but not limited to superoxide (Kalyanaraman et al., 2017), was elevated in mtD2⁺ macrophages compared with mtD2⁻ macrophages (Figure 3G). Second, we exposed WAT SVF isolates to 1-μm red fluorescent latex beads and found that mtD2⁺ macrophages exhibited significantly decreased bead phagocytosis than mtD2⁻ macrophages from paired samples (Figure 3H).

Next, we sort-purified mtD2⁻ and mtD2⁺ macrophages from eWAT of MitoFat mice and performed unbiased RNA sequencing (Table S1). Principal component analyses indicated that eWAT macrophages that had internalized mitochondria from adipocytes were transcriptionally distinct from macrophages that had not taken up mitochondria (Figure 3I). Differential gene expression analyses showed that 120 genes were significantly downregulated, and 219 genes were significantly upregulated at least 2-fold (Figure 3J). The top 20 differentially expressed genes in mtD2⁺ versus mtD2⁻ macrophages included upregulated mtDNA-encoded, chemokine, and anti-inflammatory genes and downregulated major histocompatibility complex (MHC)-II antigen presentation genes (Figure S2). Further, gene set enrichment analysis (GSEA) indicated that macrophages that had received mitochondria from adipocytes were enriched in genes associated with the hypoxia-inducible factor (HIF)-1α/tissue factor (TF) pathway (Figure 3K) and de-enriched for genes associated with electron transport (Figure 3L) and collagen synthesis (Figure 3M). Collectively, these data indicate that




Figure 2. Intravital 2-Photon Microscopy Reveals Mitochondria Transfer in WAT In Vivo

(A) Experimental design. (B) Low power magnification showing host adipocytes (dashed circular line), host-derived mtD2⁺ mitochondria (green, arrows), and a WT CMTMR-labelled macrophage (red). Scale bar, 10 µm. (C) High-power magnification Z stack of donor macrophage, showing yellow host-derived mitochondria within a donor macrophage (mac). Scale bar, 5 µm. (D) Three-dimensional reconstruction and positional mapping of host cell-derived mtD2⁺ mitochondria in a donor macrophage. Horizontal and vertical lines show coordinates. (E) Time-lapse imaging revealing a mitochondria uptake event in WAT *in vivo*. Representative of n = 4 mice from 2 independent experiments with n = 2 mtD2 controls without adoptive transfer.

mitochondria transfer from adipocytes to macrophages defines a distinct macrophage subpopulation.

A Genome-wide CRISPR-Cas9 Knockout Screen Identifies the HS Biosynthetic Pathway as Essential for Mitochondria Uptake

Mitochondria uptake is known to be diminished by various cytochalasins, a family of fungal metabolites that inhibit actin polymerization and the ability to rearrange the cytoskeleton (Boudreau et al., 2014; Kitani et al., 2014). However, little is known about which genes are required for mitochondria uptake. N-formylmethionine residues are found only in bacterial and mtDNA-encoded proteins and bind formyl peptide receptor 1 (FPR1), a gene that is highly expressed by macrophages (Gemperle et al., 2012; He and Ye, 2017). Therefore, we hypothesized that FPR1 might mediate mitochondria transfer from adipocytes to macrophages. To test this, we crossed MitoFat to *Fpr1*^{-/-} mice to generate FPR1-deficient MitoFat mice and, unexpectedly, found that FPR1 is not required for mitochondria transfer from adipocytes to macrophages *in vivo* (Figure S3). While we cannot exclude a role for other FPR family members in mediating mitochondria transfer, we turned to an unbiased approach to provide further insight into the molecular mechanisms mediating mitochondria uptake by macrophages.

We employed a genome-wide CRISPR-Cas9 knockout screen to identify genes that are essential for mitochondria uptake. To accomplish this, we stably transduced BV2 myeloid cells to express Cas9, introduced the Brie CRISPR library (Doench et al., 2016) with ~4 sgRNA guides targeting each of 19,674 genes (Doench et al., 2016; Orvedahl et al., 2019), and then exposed the pool of cells to mtD2 mitochondria (Figures 4A, S4A, and S4B). We then sort-purified mtD2⁻ BV2 cells that failed to physically interact with exogenous mitochondria (Figure S4B). The sgRNA guides were amplified and sequenced in the mtD2⁻ and input populations, and STARS score enrichment analysis was

performed. This screen identified 23 candidate genes with a false discovery rate (FDR) <0.05, each with a STARS score >4.5 (Figure 4B; Table S2). Of these 23 genes, 13 are involved in HS biosynthesis (Lindahl et al., 2015) (red genes, Figure 4C), which was confirmed by gene ontology (GO) term enrichment analysis that indicated a >200-fold enrichment for genes related to HS proteoglycan (HSPG) biosynthetic process (Figure 4D). These 13 genes traced the entire HS biosynthetic process (Lindahl et al., 2015), including the genes required for synthesis of HS substrates (*Uxs1*, *Ugp2*, and *Ugdh*), the glycosaminoglycan (GAG) linker chain (*Fam20b*, *B4galt7*, *B3galt6*, and *B3gat3*), and the HS chain (*Extl3*, *Ext1*, and *Ext2*), as well as genes involved in HS polymer sulfation (*Papss1*, *Slc35b2*, and *Hs6st1*) (Figure 4E).

To validate that HSs are required for optimal mitochondria uptake, we used CRISPR sgRNA guides directed against *B3gat3*, *B3galt6*, *B4galt7*, *Uxs1*, *Ugp2*, *Ugdh*, *Extl3*, *Ext1*, *Ext2*, *Slc35b2*, *Papss1*, and *Hs6st1* to generate gene-edited clonal cell lines. As a control, we used a CRISPR sgRNA guide against green fluorescent protein (GFP). Each gene-edited cell line had undetectable HS on their cell surface, except for sgRNA *Hs6st1*, a cell line that is capable of synthesizing intact HS chains but which lacks 6-O sulfation (Figures 4F and S4C). Each gene-edited cell line exhibited significantly decreased mitochondria uptake compared with control cells (Figure 4G). In contrast, deletion of these HS synthesis genes did not affect uptake of 1-µm latex beads (Figure 4H), suggesting that HS can facilitate interactions with mitochondria but not all foreign objects. Further supporting a role for HS in mitochondria uptake, we found that enzymatic removal of HS on BV2 cells with heparanases I-III impaired mitochondria uptake (Figure 4I). Consistent with these data, pre-treating purified mitochondria with HS proteoglycan (HSPG) inhibited their uptake by BV2 cells *in vitro* (Figure 4J). In addition, treating MitoFat mice with unfractionated heparin, a highly sulfated form of HS that is widely used clinically as an anticoagulant (Oduah et al., 2016), inhibited mitochondria transfer from adipocytes to macrophages in eWAT *in vivo* (Figure 4K). These genetic, enzymatic, and pharmacological approaches




Figure 3. Adipocytes Transfer Mitochondria to Macrophages in WAT *In Vivo*, Defining a Distinct Macrophage Subpopulation

(A and B) (A) Frequencies and (B) numbers of live CD45⁺ CD11b⁺ CD64⁺ macrophages per gram of eWAT from control (Con, n = 9) and adipocyte-specific mitochondria reporter mice (MitoFat, n = 7). (C) WT eWAT macrophages (mac) were labeled with CMTMR and adoptively transfer into MitoFat mice for intravital 2-photon microscopy of host eWAT. Three-dimensional reconstruction (left) and positional mapping (right) of donor macrophage containing adipocyte-derived mitochondria (arrows). (D–G) (D) Gating strategy to define macrophages that have (mtD2⁺) or have not (mtD2[−]) acquired mitochondria from adipocytes *in vivo* in MitoFat mice (n = 5/group) for comparison of fluorescence intensity of (E) the charge-independent mitochondrial mass indicator MitD-Red, (F) the charge-dependent mitochondria dye MitoTracker Red CMXRos (CMXRos) normalized to MitD-Red, (G) the charge-dependent dye MitSOX Red normalized to CMXRos.

(H) Comparison of mtD2⁺ and mtD2[−] eWAT macrophage phagocytosis of 1-μm polyred latex beads (n = 6/group).

For (E)–(H), lines connect paired data points obtained from the same sample.

(I and J) Messenger RNA sequencing of mtD2⁺ and mtD2[−] macrophages from eWAT of MitoFat mice (n = 4).

(I) Principal component analysis with dashed lines connecting paired samples from the same mouse and shaded ellipses defining 95% confidence intervals for each group.

(J) Volcano plot of detected transcripts.

(K–M) GSEA showing enrichment of the HIF-1α/TF pathway (K) and de-enrichment of genes associated with electron transport (L) and collagen synthesis (M). Data expressed as mean ± SEM. Student's t test (A and B) and paired t test (E–H); *p < 0.05, ***p < 0.001. See also Figure S2.

collectively indicate that HS are essential for efficient mitochondria uptake both *in vitro* and *in vivo*.

Obesity Is Associated with Decreased Mitochondria Transfer from Adipocytes to Macrophages

To test whether intercellular mitochondria transfer from adipocytes to macrophages is altered in the setting of obesity, we fed MitoFat mice a normal chow or HFD for 12 weeks. HFD-induced obesity was associated with a marked decrease in intercellular mitochondria transfer from adipocytes to macrophages in WAT (Figures 5A and 5B). To test whether this obesity-associated change is due to impaired release of mitochondria from adipocytes, impaired uptake of mitochondria by macrophages, or both, we utilized a WAT explant co-culture system. Specifically, minced WAT explants from CD45.1 mtD2 mice fed a chow or HFD for 10–12 weeks were co-cultured with minced WAT explants from CD45.2 WT mice fed a chow or HFD for 10–12 weeks. Flow cytometry was used to determine the proportion of CD45.2 WT macrophages that had received mitochondria from mtD2 donor WAT. We found that varying the nutritional status of the donor mtD2 WAT had no effect on mitochondria transfer to recipient macrophages (Figure 5C). In contrast, we observed significantly decreased mitochondria transfer when the recipient macrophages were from HFD WAT (Figure 5C). Thus, the decrease in mitochondria transfer from adipocytes to macrophages appears

to be due to a macrophage-intrinsic impairment in mitochondria uptake. Consistent with this result, we observed that eWAT macrophages from HFD-fed mice have substantially decreased levels of HS on their surface compared with normal chow-fed controls (Figure 5D).

In obesity, pro-inflammatory cytokines such as IFN-γ, LPS, and other stimuli promote the accumulation of M1-like macrophages in WAT (Brestoff and Artis, 2015; Caesar et al., 2012; Cani et al., 2007; Clemente-Postigo et al., 2019; Hotamisligil, 2017; Lee et al., 2018). We found that the proportion of F4/80⁺ CD64⁺ CD206[−] CD11c⁺ (M1-like) macrophages that had received mitochondria from adipocytes was significantly lower than in F4/80⁺ CD64⁺ CD206⁺ CD11c[−] (M2-like) macrophages in mice fed a chow diet (Figure 5E). Further, both M1-like and M2-like WAT macrophage populations exhibited a decrease in the proportion of mtD2⁺ cells in the setting of HFD-induced obesity (Figure 5E), and the percentage of mtD2⁺ macrophages correlated with cell surface HS levels (Figure 5F). Therefore, we hypothesized that M1-like polarization with IFN-γ and LPS might impair mitochondria uptake. To test this idea, we polarized BV2 cells toward an M0, M2, or M1 phenotype with phosphate-buffered saline (PBS), IL-4, or IFN-γ plus LPS, respectively, for 24 h. Polarization was confirmed by intracellular flow cytometric analysis of the M2 marker Arginase 1 (ARG1, Figure S5A) and the M1 marker Inducible nitric oxide synthase (iNOS) (Figure S5B). We




Figure 4. Genome-wide CRISPR Knockout Screen Identifies HSs as Essential for Mitochondria Uptake by Macrophages

(A) Experimental design of genome-wide CRISPR-Cas9 knockout screen.

(B) STARS score analyses identifies 23 enriched genes with FDR < 0.05.

(C) List of 23 enriched genes, of which red genes are required for HS biosynthesis.

(D) GO Term enrichment analysis of the 23 genes identified by the CRISPR screen.

(E) Schematic relating the 13 HS synthesis genes enriched in the CRISPR screen, tracing the entire HS biosynthesis pathway unsupervised.

(F–H) (F) Surface HS levels, (G) mtD2 mitochondria uptake, and (H) 1-μm red latex bead phagocytosis in 13 clonally selected BV2-Cas9 cell lines, with sgRNA targeting the indicated genes. n = 3 independent experiments, one-way ANOVA with Dunnett post-hoc test comparing against the sgRNA to *Gfp* control; *p < 0.05, **p < 0.01, ***p < 0.001.

(I) Mitochondria uptake after pre-treating BV2 cells with heparanases I–III (5 U/mL each), n = 3/group.

(J) Mitochondria uptake by BV2 cells after pre-incubation of mtD2 mitochondria with 1 mg/mL HS proteoglycan (HSPG), n = 4/group.

(K) Frequencies of mtD2⁺ macrophages in eWAT from MitoFat mice after 7 days of treatment with PBS or heparin (5 mg/kg, n = 3/group).

For (I)–(K), Student's t test; ***p < 0.001. *p < 0.05, **p < 0.01, ***p < 0.001. For (F)–(K), data are expressed as mean ± SEM. See also Figure S4.

then cultured M0-, M1-, and M2-polarized BV2 cells with 1-μm red latex beads, which are similar in size to mitochondria, and found that IFN-γ and LPS treatment resulted in increased bead phagocytosis compared with IL-4 (Figures 5G and S5C). This observation is consistent with prior studies showing that activation of macrophages with IFN-γ increases both phagocytosis and macropinocytosis (Bosedasgupta and Pieters, 2014). In contrast, treatment of BV2 cells with IFN-γ plus LPS markedly decreased the uptake of mitochondria, whereas IL-4 had no effect (Figures 5H and S5D). We also found that IFN-γ plus LPS decreased the expression of transcripts involved in HS biosynthesis, whereas IL-4 had no effect (Figure 5I). These data indicate that IFN-γ plus LPS activation impairs mitochondria uptake and suggest that these pro-inflammatory stimuli may contribute to the reduction in adipocyte-to-macrophage mitochondria transfer observed in murine obesity.

Genetic Deletion of *Ext1* from Myeloid Cells Impairs Mitochondria Transfer to Macrophages and Promotes Fat Accumulation

To investigate the role of intercellular mitochondria transfer in regulating systemic metabolism, we conditionally deleted *Ext1* from myeloid cells using *Lyz2-Cre*. Deletion of *Ext1* in macrophages did not affect the frequencies (Figure 6A) and numbers (Figure 6B) of macrophages in eWAT. This indicates that *Ext1* is dispensable for maintaining macrophage abundance in chow-fed mice. Next, we confirmed that eWAT macrophages from *Ext1*^{ΔLyz2} mice had reduced levels of HS on their surface (Figure 6C) and impaired mitochondria uptake (Figure 6D)

compared with *Ext1*^{F/F} littermate controls. Moreover, we found that *Ext1*^{ΔLyz2} mice had significantly higher body weight (Figure 6E), absolute eWAT mass (Figure 6F), and relative eWAT mass normalized to body weight (Figure 6G) compared with *Ext1*^{F/F} littermate controls. Whole-body lean mass did not differ between *Ext1*^{ΔLyz2} and *Ext1*^{F/F} groups; however, whole-body fat mass (Figure 6H) and adiposity (Figure 6I) were significantly increased in *Ext1*^{ΔLyz2} mice. These changes were accompanied by impaired glucose tolerance (Figure 6J) and insulin tolerance (Figure 6K) in *Ext1*^{ΔLyz2} mice compared with controls.

These observations suggested that *Ext1*^{ΔLyz2} mice have altered energy homeostasis. To investigate this hypothesis, we performed metabolic cage analyses and found that *Ext1*^{ΔLyz2} mice exhibited significantly lower energy expenditure than *Ext1*^{F/F} littermates on a chow diet (Figures 7A and 7B). The respiratory quotient (RQ) did not differ between the two groups (Figure 7C), indicating that energy substrate utilization may be unaltered in *Ext1*^{ΔLyz2} mice. Furthermore, food intake (Figure 7D) and physical activity (Figure 7E) also did not differ between the two groups. Given these findings, we hypothesized that *Ext1*^{ΔLyz2} mice would exhibit more severe diet-induced obesity. Indeed, we found that *Ext1*^{ΔLyz2} mice exhibited increased body weight (Figure 7F) as well as eWAT and iWAT masses (Figure 7G) following 8 weeks of HFD feeding compared with *Ext1*^{F/F} littermates. BAT masses did not differ significantly between groups (Figure 7G). In addition, HFD feeding resulted in more severe glucose intolerance in *Ext1*^{ΔLyz2} mice compared with *Ext1*^{F/F} controls (Figure 7H). Taken together, these data indicate that conditional deletion of *Ext1* in myeloid cells impairs mitochondria




Figure 5. Intercellular Mitochondria Transfer to Macrophages Is Impaired in Obesity and Following Stimulation with IFN- γ and LPS

(A and B) Male 6-week-old MitoFat mice were fed a chow or HFD for 12 weeks, and live CD45⁺ SiglecF⁺ CD11b⁺ CD64⁺ macrophages in eWAT macrophages were identified by flow cytometry. The proportion of mtD2⁺ macrophages are shown. $n = 6$ –8/group from 3 independent experiments.

(C) Frequency of WT recipient macrophages that are mtD2⁺ after WAT explant co-culture with congenically disparate mtD2 donor WAT. C, chow diet; H, high-fat diet; $n = 5$ /group from 2 independent experiments combined.

(D) Relative surface HS levels on eWAT macrophages in mice fed a chow or HFD for 10–12 weeks, representative of $n = 6$ –8 mice/group pooled from 2 independent experiments.

For (A)–(D), Student's t test; *** $p < 0.001$.

(E) Frequencies of mtD2⁺ cells in CD206⁺ CD11c[−] and CD206[−] CD11c⁺ macrophages from eWAT of MitoFat mice fed a chow ($n = 4$) or HFD ($n = 3$) for 12 weeks.

(F) Pearson linear regression correlating the proportion of mtD2⁺ eWAT macrophages defined in (E) to their relative HS levels. Gray, chow; black, HFD.

(G–I) BV2 cells were stimulated with PBS, 20 ng/mL interleukin (IL-4), or 10 ng/mL interferon (IFN)- γ plus 1 ng/mL lipopolysaccharide (LPS) for 24 h before co-culturing with 1- μ m polyed latex beads and purified mtD2 mitochondria. Frequencies of cells that took up (G) beads and (H) mitochondria. (I) mRNA expression of the indicated HS biosynthesis genes in BV2 cells. $n = 4$ independent experiments, one-way ANOVA with Fisher's LSD post-hoc test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Data are represented as mean \pm SEM. See also Figure S5.

transfer and decreases energy expenditure, promoting the accumulation of fat mass.

DISCUSSION

Here, we provide evidence that adipocytes transfer their mitochondria to macrophages in WAT *in vivo*, and that this process defines a transcriptionally distinct macrophage subpopulation. We also demonstrate that this process is decreased in the context of diet-induced obesity due to reduced mitochondria uptake by WAT macrophages. Mitochondria uptake is mediated by HS, and surface levels of HS are decreased on WAT macrophages from obese mice and in BV2 cells following stimulation with IFN- γ and LPS. Furthermore, conditional deletion of EXT1 in myeloid cells reduces WAT macrophage HS levels, impairs mitochondria uptake, promotes fat mass accumulation, and reduces systemic energy expenditure without affecting food intake or physical activity levels.

Previous studies have demonstrated that mitochondria are released by various cell types *in vitro* and *in vivo* and that functional mitochondria circulate in murine and human peripheral blood (Al Amir Dache et al., 2020; Boudreau et al., 2014; Chiu et al., 2003; Maeda and Fadeel, 2014; Pollara et al., 2018; Scozzi

et al., 2019; Torralba et al., 2016). It also has been reported that mitochondria-deficient cell types can capture purified mitochondria from their environment and utilize them to support cellular metabolic demands and cell proliferation (Kim et al., 2018; Kitani et al., 2014; Spees et al., 2006). *In vivo*, cancer cells including a mitochondria-deficient ρ^0 melanoma cell line can acquire mitochondria from host cells (Dong et al., 2017; Rebbeck et al., 2011), and ischemic neurons acquire mitochondria from astrocytes in the brain in a murine model of ischemic stroke (Haya-kawa et al., 2016). These and other observations have led to the hypothesis that intercellular mitochondria transfer may serve as a mechanism to rescue or sustain metabolically compromised cells (Torralba et al., 2016). In our studies, macrophages that have acquired mitochondria from adipocytes appear to produce more mitochondrial ROS and exhibit evidence of hypoxia and de-enrichment of nuclear-encoded genes involved in mitochondrial homeostasis and maintenance of the electron transport chain. Although deletion of *Ext1* from myeloid cells impaired mitochondria uptake by WAT macrophages without affecting macrophage abundance at steady state, further research is needed to determine whether and how mitochondria transfer affects nuclear gene expression and supports the metabolic demands of macrophages.




Figure 6. Genetic Deletion of *Ext1* in Myeloid Cells Impairs Mitochondria Transfer to Macrophages and Is Associated with Increased Fat Mass and Glucose Intolerance

(A and B) (A) Frequencies and (B) numbers per gram of eWAT macrophages in *Ext1*^{ΔLyz2} mice (n = 15) and *Ext1*^{F/F} littermate controls (n = 14) at steady state. Pre-gated on live CD45⁺ SiglecF[−] cells.

(C) Relative surface HS levels on eWAT macrophages in *Ext1*^{ΔLyz2} mice (n = 6) and *Ext1*^{F/F} littermate controls (n = 6).

(D) Frequencies of *Ext1*^{ΔLyz2} (n = 7) and *Ext1*^{F/F} (n = 8) eWAT recipient macrophages that are mtD2+ after eWAT explant co-culture with CD45.1 mtD2 donor WAT.

(E–G) (E) Body weight, (F) absolute eWAT mass, and (G) relative eWAT mass normalized to body weight in *Ext1*^{ΔLyz2} mice (n = 21) and *Ext1*^{F/F} littermate controls (n = 24).

(H–K) (H) Whole-body lean and fat masses, (I) adiposity, (J) glucose tolerance tests, and (K) insulin tolerance tests in 5–6-month-old *Ext1*^{ΔLyz2} mice (n = 8) and *Ext1*^{F/F} littermate controls (n = 11).

For (A)–(I), Student's t test. For (J) and (K), two-way ANOVA with repeated measures with Fisher's LSD post hoc test. *p < 0.05, **p < 0.01, ***p < 0.001. Data are represented as mean ± SEM.

Emerging evidence supports a role for intercellular mitochondria transfer in regulating the immune system. For example, the capture of exogenous mitochondria is reported to promote regulatory T cell programming at steady state, suggesting that mitochondria transfer might have anti-inflammatory properties (Court et al., 2020). In contrast, mitochondria capture by neutrophils promotes pulmonary inflammation following lung transplantation (Scozzi et al., 2019). Our studies indicate that mitochondria uptake by macrophages occurs in WAT frequently in healthy conditions when macrophages are polarized toward an M2-like activation state driven by type 2 cytokines, such as IL-4 and IL-13, which are produced by group 2 innate lymphoid cells, eosinophils, and other cell types present in healthy WAT (Brestoff and Artis, 2015; Lee et al., 2018). In comparison, the frequency of mitochondria transfer from adipocytes to macrophages is markedly decreased in obesity, a pro-inflammatory state in which macrophages are exposed to factors that induce a type 1 im-

mune response, including but not limited to IFN-γ and LPS (Brestoff and Artis, 2015; Caesar et al., 2012; Cani et al., 2007; Clemente-Postigo et al., 2019; Hotamisligil, 2017; Makki et al., 2013; Reilly and Saltiel, 2017). This suggests that M1-like polarization inhibits mitochondria uptake. Consistent with this, the frequency of mitochondria transfer from adipocytes to CD206⁺ CD11c[−] (M2-like) macrophages is higher than to CD206[−] CD11c⁺ (M1-like) macrophages, and HFD-induced obesity is associated with decreased mitochondria transfer from adipocytes to both macrophage populations. Further, BV2 cells also exhibit decreased mitochondria uptake following IFN-γ and LPS stimulation. These observations implicate a reduction in mitochondria transfer to be a feature of metabolic diseases such as obesity, at least in WAT.

The mechanisms by which mitochondria are taken up into cells are not well understood. Mitochondria uptake is inhibited by cytochalasins, which impair actin cytoskeleton




Figure 7. Deletion of *Ext1* in Myeloid Cells Is Associated with Decreased Energy Expenditure and Increased Susceptibility to Diet-Induced Obesity

(A–E) Metabolic cage analyses of *Ext1^{ΔLyz2}* mice ($n = 19$) and *Ext1^{F/F}* littermate controls ($n = 17$). Energy expenditure expressed as heat produced per kg/body weight per hour at (A) 5.5-min resolution and (B) on average for light versus dark phases. (C) RQ during the light versus dark phases. (D) Food intake and (E) physical activity levels.

(F–H) (F) Body weight, (G) indicated adipose tissue depot masses, and (H) glucose tolerance tests of 5–6-month-old *Ext1^{ΔLyz2}* mice ($n = 14$) and *Ext1^{F/F}* littermate controls ($n = 12$) fed a HFD for 9 weeks.

For (A)–(C) and (H), two-way ANOVA with repeated measures and LSD post-hoc test. For (D)–(F), Student's *t* test. For (G), two-way ANOVA with LSD post hoc test. * $p < 0.05$, ** $p < 0.01$. Data are represented as mean \pm SEM.

rearrangement (Boudreau et al., 2014; Kitani et al., 2014). However, there is no known receptor for intact mitochondria. Our genome-wide CRISPR-Cas9 knockout screen identified 23 candidate genes that might contribute to mitochondria uptake. Of these, 13 genes traced the HS biosynthesis pathway in an unsupervised analysis, strongly implicating HS as an attachment factor to support mitochondria uptake. Deletion of these HS biosynthesis genes impaired mitochondria uptake by BV2 cells *in vitro* without affecting phagocytosis of latex beads of a similar size as mitochondria. This suggests that HS species may confer some specificity to bind mitochondria but not to all foreign objects. Further supporting a role for HS in mitochondria uptake, stimulating BV2 cells with IFN- γ and LPS downregulated expression of *Ext2* and *Hs6st1*, genes that are required for HS synthesis and 6-O sulfation, respectively, and eWAT macrophages from HFD-induced obese mice had lower surface HS levels than chow-diet-fed controls. The degree of mitochondria transfer from adipocytes to macrophages is correlated with macrophage HS levels. In addition, heparin treatment was sufficient to inhibit mitochondria transfer from adipocytes to macrophages *in vivo*, and conditional deletion of *Ext1* from myeloid cells in mice reduced eWAT macrophage HS levels and mitochondria uptake by WAT macrophages. These data support a role for HS in mediating mitochondria uptake by murine macrophages *in vitro* and *in vivo*. These observations raise questions about whether treating patients with heparin for anti-coagulation results in impaired intercellular mitochondria transfer. Heparin is known to cause metabolic derangements (Lee et al., 1988; Rutstein et al.,

1969; Zhu et al., 2017) and reduce inflammation (Ekre et al., 1992; Mousavi et al., 2015) in mice and humans, and further research is needed to determine whether intercellular mitochondria transfer contributes to these effects. Further research will also be needed to determine whether HS mediates mitochondria transfer or obesity pathogenesis in humans and whether other factors or genes are also required for mitochondria uptake or transfer.

The generation of *Ext1^{ΔLyz2}* mice allowed us to investigate whether the genetic disruption of mitochondria transfer affects macrophage abundance or systemic metabolism. We found that mice lacking *Ext1* in myeloid cells had unaltered macrophage frequencies or numbers in WAT, suggesting that HS and mitochondria transfer is not required to sustain WAT macrophage abundance. However, *Ext1^{ΔLyz2}* mice exhibited mildly increased body weight and an approximately 30% increase in absolute and relative eWAT mass relative to littermate controls. Metabolic cage analyses indicated that conditional deletion of *Ext1* in myeloid cells resulted in substantially lower energy expenditure without affecting food intake or physical activity, establishing positive energy balance that favors accumulation of fat mass over time. Thus, it appears that intercellular mitochondria transfer to macrophages might contribute to the maintenance of energy homeostasis in mice. These data further raise the possibility that the obesity-associated decrease in mitochondria transfer from adipocytes to macrophages might have a role in weight gain or the metabolic dysfunction in response to an HFD. Additional research will be needed to understand how

mitochondria transfer affects macrophage function and WAT and BAT homeostasis.

Interactions between adipocytes and immune cells are essential to maintain metabolic homeostasis and, when dysregulated, can drive pathologic inflammation that promotes obesity and obesity-associated metabolic dysfunction (Brestoff and Artis, 2015; Lee et al., 2018; Singer and Lumeng, 2017; Sun et al., 2011). Here, we identified that intercellular mitochondria transfer occurs *in vivo* in WAT and defined an adipocyte-to-macrophage mitochondria transfer axis that is dysregulated in obesity due to classical or M1-like macrophage polarization. Intercellular mitochondria transfer occurs, at least in part, through a HS-dependent mechanism both *in vitro* and *in vivo*, and genetic disruption of HS synthesis on macrophages impairs intercellular mitochondria transfer and compromises energy homeostasis, raising the possibility of a functional relationship. Indeed, patients and mice with heterozygous loss-of-function mutations in the HS synthesis gene *Ext1* have impaired lipid and glucose homeostasis (Mooij et al., 2015). Furthermore, genome-wide association studies have linked single nucleotide polymorphisms in *Ext2*, which forms a heterodimer with EXT1 to polymerize the HS chain (Busse et al., 2007), to increased risk for type 2 diabetes (Sladek et al., 2007). Thus, our findings reveal a new paradigm of immunometabolic crosstalk in which some cells, such as adipocytes, transfer their mitochondria to macrophages to regulate systemic metabolic homeostasis. These observations suggest that intercellular mitochondria transfer may have evolved as a homeostatic process that enables immune cells to respond to and regulate their local tissue microenvironment. Therapeutically targeting intercellular mitochondria transfer may represent an unappreciated strategy to ameliorate metabolic diseases.

Limitations of Study

A significant limitation of these studies is that they were done in mice and cells isolated from mice, not humans. Improved tools and techniques to study intercellular mitochondria transfer in humans, especially *in vivo*, are needed to determine the translational relevance of intercellular mitochondria transfer in the context of human metabolic homeostasis, immunity, obesity pathogenesis, or other disease states. Furthermore, it remains unknown whether intercellular mitochondria transfer to macrophages affects the bioenergetic or metabolic status of the recipient cell, a topic that warrants further study. Although we identify a role for HS in mediating intercellular mitochondria transfer *in vitro* and *in vivo* in mice, targeting the HS pathway with pharmacologic, enzymatic, and genetic approaches resulted in a partial disruption in mitochondria uptake. This may have impacted the severity of the metabolic abnormalities observed in mice with conditional deletion of *Ext1* in myeloid cells and suggests that there are likely other factors and/or genes that also facilitate the mitochondria uptake process.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

● KEY RESOURCES TABLE

● RESOURCE AVAILABILITY

- Lead Contact
- Materials Availability
- Data and Code Availability

● EXPERIMENTAL MODEL AND SUBJECT DETAILS

- Mouse Models
- Cell Culture

● METHOD DETAILS

- Bone Marrow Chimeras
- Metabolic Cage Analyses
- Heparin Treatment
- Glucose Homeostasis Analyses
- Body Composition Analysis
- Isolation of Immune Cells from Mouse Tissues
- Flow Cytometry and Cell-Sorting
- Staining eWAT Macrophages with Mitochondria Dyes
- Adoptive Cell Transfer
- Intravital 2-Photon Microscopy
- RNA Sequencing and Analysis
- Mitochondria and Bead Uptake Assays in BV2 Cells
- Quantitative Real-Time PCR
- Mitochondria Isolation
- Brie CRISPR Library Preparation
- Genome-Wide CRISPR Knockout Screen, Sequencing, and Analysis
- Generation of BV2 Knockout Cells
- WAT Co-culture System

● QUANTIFICATION AND STATISTICAL ANALYSIS

- Statistical Analyses

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.cmet.2020.11.008>.

ACKNOWLEDGMENTS

This work was supported by National Institutes of Health (NIH) R01-DK111389, NIH Merit award R37-AR046523, Shriner's Hospital for Children grant #85400, and a grant from Siteman Cancer Center to S.L.T. Additional funding was provided by the NIH Director's Early Independence Award (DP5 OD028125), the Burroughs Wellcome Fund (CAMS 1019648), and Children's Discovery Institute (MI-F-2019-795) to J.R.B. Support was also provided by NIH R01-AI123348 (M.S.D.), U19-AI142784 (H.W.V.), K08-AR065577 (B.S.K.), R01-AR070116 (B.S.K.), Doris Duke Charitable Foundation (B.S.K.), Celgene Corporation (B.S.K.), Leo Pharma (B.S.K.), and K99-HL138163 (J.W.W.). The Genome Technology Access Center (GTAC) is partially supported by NCI Cancer Center Support grant P30 CA91842 to Siteman Cancer Center, ICTS/CTSA grant# UL1TR002345 from the National Center for Research Resources (NCRR), a component of the National Institutes of Health (NIH), and NIH Roadmap for Medical Research. We thank D. Brinja and E. Lantelme at the flow cytometry and cell sorting Core and E. Tycksen at GTAC at Washington University School of Medicine for their technical support.

AUTHOR CONTRIBUTIONS

J.R.B., C.B.W., J.R.M., M.N.R., N.P.M., Y.L., W.Z., H.M., B.T.S., M.R.M., B.L.S., D.R.B., A.O., and J.W.W. performed experiments. J.R.B., C.B.W., J.R.M., H.M., D.B., A.O., B.H.Z., S.A.H., B.S.K., J.D., G.J.R., M.S.D., H.W.V., A.E.G., and S.L.T. designed the project. J.R.B., C.B.W., J.R.M., Y.L., B.T.S., M.R.M., B.L.H. Jr., W.Z., N.R., X.W., B.H.Z., and J.W.W. analyzed the data. J.R.B. wrote the paper. All authors read, edited, and approved the manuscript.

DECLARATION OF INTERESTS

D.R.B. and H.W.V. are employees of J. Virol. Biotechnol., a for-profit institution. H.W.V. is a founder of PierenianDx and Casma Therapeutics. M.S.D. is a consultant for Inbios and on the Scientific Advisory Board of Moderna. B.S.K. has served as a consultant for AbbVie, Inc., Concert Pharmaceuticals, Incyte Corporation, Menlo Therapeutics, and Pfizer, Inc; has participated on the advisory board for Celgene Corporation, Kiniksa Pharmaceuticals, Menlo Therapeutics, Regeneron Pharmaceuticals, Inc., Sanofi, and Theravance Pharmaceuticals; is a stockholder of Gilead Sciences, Inc. and Mallinckrodt Pharmaceuticals; and is a Founder and Chief Scientific Officer of Nuogen Pharma, Inc. The other authors declare no competing interests.

Received: April 22, 2020

Revised: September 3, 2020

Accepted: November 11, 2020

Published: December 4, 2020

REFERENCES

- Al Amir Dache, Z., Otandault, A., Tanos, R., Pastor, B., Meddeb, R., Sanchez, C., Arena, G., Lasorsa, L., Bennett, A., Grange, T., et al. (2020). Blood contains circulating cell-free respiratory competent mitochondria. *FASEB J.* **34**, 3616–3630.
- Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., et al. (2000). Gene ontology: tool for the unification of biology. *Nat. Genet.* **25**, 25–29.
- Bosedasgupta, S., and Pieters, J. (2014). Inflammatory stimuli reprogram macrophage phagocytosis to macropinocytosis for the rapid elimination of pathogens. *PLoS Pathog.* **10**, e1003879.
- Boudreau, L.H., Duchez, A.C., Cloutier, N., Soulet, D., Martin, N., Bollinger, J., Paré, A., Rousseau, M., Naika, G.S., Lévesque, T., et al. (2014). Platelets release mitochondria serving as substrate for bactericidal group IIA-secreted phospholipase A2 to promote inflammation. *Blood* **124**, 2173–2183.
- Brestoff, J.R., and Artis, D. (2015). Immune regulation of metabolic homeostasis in health and disease. *Cell* **161**, 146–160.
- Brestoff, J.R., Kim, B.S., Saenz, S.A., Stine, R.R., Monticelli, L.A., Sonnenberg, G.F., Thome, J.J., Farber, D.L., Lutfy, K., Seale, P., and Artis, D. (2015). Group 2 innate lymphoid cells promote beiging of white adipose tissue and limit obesity. *Nature* **519**, 242–246.
- Busse, M., Feta, A., Presto, J., Wilén, M., Grønning, M., Kjellén, L., and Kusche-Gullberg, M. (2007). Contribution of EXT1, EXT2, and EXTL3 to heparan sulfate chain elongation. *J. Biol. Chem.* **282**, 32802–32810.
- Caesar, R., Reigstad, C.S., Bäckhed, H.K., Reinhardt, C., Ketonen, M., Lundén, G.O., Cani, P.D., and Bäckhed, F. (2012). Gut-derived lipopolysaccharide augments adipose macrophage accumulation but is not essential for impaired glucose or insulin tolerance in mice. *Gut* **61**, 1701–1707.
- Camell, C.D., Sander, J., Spadaro, O., Lee, A., Nguyen, K.Y., Wing, A., Goldberg, E.L., Youm, Y.H., Brown, C.W., Elsworth, J., et al. (2017). Inflammasome-driven catecholamine catabolism in macrophages blunts lipolysis during ageing. *Nature* **550**, 119–123.
- Cani, P.D., Amar, J., Iglesias, M.A., Poggi, M., Knauf, C., Bastelica, D., Neyrinck, A.M., Fava, F., Tuohy, K.M., Chabo, C., et al. (2007). Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes* **56**, 1761–1772.
- Chang, J.C., Chang, H.S., Wu, Y.C., Cheng, W.L., Lin, T.T., Chang, H.J., Kuo, S.J., Chen, S.T., and Liu, C.S. (2019). Mitochondrial transplantation regulates antitumour activity, chemoresistance and mitochondrial dynamics in breast cancer. *J. Exp. Clin. Cancer Res.* **38**, 30.
- Chiu, R.W., Chan, L.Y., Lam, N.Y., Tsui, N.B., Ng, E.K., Rainer, T.H., and Lo, Y.M. (2003). Quantitative analysis of circulating mitochondrial DNA in plasma. *Clin. Chem.* **49**, 719–726.
- Clemente-Postigo, M., Oliva-Olivera, W., Coin-Aragüez, L., Ramos-Molina, B., Giraldez-Perez, R.M., Lhamyani, S., Alcaide-Torres, J., Perez-Martinez, P., El Bekay, R., Cardona, F., and Tinahones, F.J. (2019). Metabolic endotoxemia promotes adipose dysfunction and inflammation in human obesity. *Am. J. Physiol. Endocrinol. Metab.* **316**, E319–E332.
- Court, A.C., Le-Gatt, A., Luz-Crawford, P., Parra, E., Aliaga-Tobar, V., Bätz, L.F., Contreras, R.A., Ortúzar, M.I., Kurte, M., Elizondo-Vega, R., et al. (2020). Mitochondrial transfer from MSCs to T cells induces Treg differentiation and restricts inflammatory response. *EMBO Rep.* **21**, e48052.
- Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21.
- Doench, J.G., Fusi, N., Sullender, M., Hegde, M., Vaimberg, E.W., Donovan, K.F., Smith, I., Tothova, Z., Wilen, C., Orchard, R., et al. (2016). Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. *Nat. Biotechnol.* **34**, 184–191.
- Dong, L.F., Kovarova, J., Bajzikova, M., Bezawork-Geleta, A., Svec, D., Endaya, B., Sachaphibulkij, K., Coelho, A.R., Sebkova, N., Ruzickova, A., et al. (2017). Horizontal transfer of whole mitochondria restores tumorigenic potential in mitochondrial DNA-deficient cancer cells. *eLife* **6**, e22187.
- Ekre, H.P., Naparstek, Y., Lider, O., Hydén, P., Hägermark, O., Nilsson, T., Vlodavsky, I., and Cohen, I. (1992). Anti-inflammatory effects of heparin and its derivatives: inhibition of complement and of lymphocyte migration. *Adv. Exp. Med. Biol.* **313**, 329–340.
- Flaherty, S.E., 3rd, Grijalva, A., Xu, X., Ables, E., Nomani, A., and Ferrante, A.W., Jr. (2019). A lipase-independent pathway of lipid release and immune modulation by adipocytes. *Science* **363**, 989–993.
- Flegal, K.M., Kruszon-Moran, D., Carroll, M.D., Fryar, C.D., and Ogden, C.L. (2016). Trends in obesity among adults in the United States, 2005 to 2014. *JAMA* **315**, 2284–2291.
- Gemperle, C., Schmid, M., Herova, M., Marti-Jaun, J., Wuest, S.J., Loretz, C., and Hersberger, M. (2012). Regulation of the formyl peptide receptor 1 (FPR1) gene in primary human macrophages. *PLoS One* **7**, e50195.
- Griessinger, E., Moschoi, R., Biondani, G., and Peyron, J.F. (2017). Mitochondrial transfer in the leukemia microenvironment. *Trends Cancer* **3**, 828–839.
- Hayakawa, K., Esposito, E., Wang, X., Terasaki, Y., Liu, Y., Xing, C., Ji, X., and Lo, E.H. (2016). Transfer of mitochondria from astrocytes to neurons after stroke. *Nature* **535**, 551–555.
- He, H.-Q., and Ye, R.D. (2017). The formyl peptide receptors: diversity of ligands and mechanism for recognition. *Molecules* **22**, 455.
- Hotamisligil, G.S. (2017). Inflammation, metaflammation and immunometabolic disorders. *Nature* **542**, 177–185.
- Islam, M.N., Das, S.R., Emin, M.T., Wei, M., Sun, L., Westphalen, K., Rowlands, D.J., Quadri, S.K., Bhattacharya, S., and Bhattacharya, J. (2012). Mitochondrial transfer from bone-marrow-derived stromal cells to pulmonary alveoli protects against acute lung injury. *Nat. Med.* **18**, 759–765.
- Jackson, M.V., and Krasnodembskaya, A.D. (2017). Analysis of mitochondrial transfer in direct co-cultures of human monocyte-derived macrophages (MDM) and mesenchymal stem cells (MSC). *Bio Protoc.* **7**, e2255.
- Jiang, D., Gao, F., Zhang, Y., Wong, D.S., Li, Q., Tse, H.F., Xu, G., Yu, Z., and Lian, Q. (2016). Mitochondrial transfer of mesenchymal stem cells effectively protects corneal epithelial cells from mitochondrial damage. *Cell Death Dis.* **7**, e2467.
- Kalyanaraman, B., Hardy, M., Podsiadly, R., Cheng, G., and Zielonka, J. (2017). Recent developments in detection of superoxide radical anion and hydrogen peroxide: opportunities, challenges, and implications in redox signaling. *Arch. Biochem. Biophys.* **617**, 38–47.
- Kim, M.J., Hwang, J.W., Yun, C.K., Lee, Y., and Choi, Y.S. (2018). Delivery of exogenous mitochondria via centrifugation enhances cellular metabolic function. *Sci. Rep.* **8**, 3330.
- Kitani, T., Kami, D., Matoba, S., and Gojo, S. (2014). Internalization of isolated functional mitochondria: involvement of macropinocytosis. *J. Cell. Mol. Med.* **18**, 1694–1703.
- Larabee, C.M., Neely, O.C., and Domingos, A.I. (2020). Obesity: a neuroimmunometabolic perspective. *Nat. Rev. Endocrinol.* **16**, 30–43.
- Lee, K.U., Lee, H.K., Koh, C.S., and Min, H.K. (1988). Artificial induction of intravascular lipolysis by lipid-heparin infusion leads to insulin resistance in man. *Diabetologia* **31**, 285–290.

- Lee, Y.S., Wollam, J., and Olefsky, J.M. (2018). An integrated view of immuno-metabolism. *Cell* 172, 22–40.
- Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 30, 923–930.
- Lin, X., Wei, G., Shi, Z., Dryer, L., Esko, J.D., Wells, D.E., and Matzuk, M.M. (2000). Disruption of gastrulation and heparan sulfate biosynthesis in EXT1-deficient mice. *Dev. Biol.* 224, 299–311.
- Lindahl, U., Couchman, J., Kimata, K., and Esko, J.D. (2015). Proteoglycans and sulfated glycosaminoglycans. In *Essentials of Glycobiology*, A. Varki, R.D. Cummings, J.D. Esko, P. Stanley, G.W. Hart, M. Aebi, A.G. Darvill, T. Kinoshita, and N.H. Packer, et al., eds. (Cold Spring Harbor), pp. 207–221.
- Liu, L., Yang, X., Wang, H., Cui, G., Xu, Y., Wang, P., Yuan, G., Wang, X., Ding, H., and Wang, D.W. (2013). Association between variants of EXT2 and type 2 diabetes: a replication and meta-analysis. *Hum. Genet.* 132, 139–145.
- Liu, R., Holik, A.Z., Su, S., Jansz, N., Chen, K., Leong, H.S., Blewitt, M.E., Asselin-Labat, M.L., Smyth, G.K., and Ritchie, M.E. (2015). Why weight? Modelling sample and observational level variability improves power in RNA-seq analyses. *Nucleic Acids Res.* 43, e97.
- Luo, W., and Brouwer, C. (2013). Pathview: an R/bioconductor package for pathway-based data integration and visualization. *Bioinformatics* 29, 1830–1831.
- Luo, W., Friedman, M.S., Shedden, K., Hankenson, K.D., and Woolf, P.J. (2009). GAGE: generally applicable gene set enrichment for pathway analysis. *BMC Bioinformatics* 10, 161.
- Maeda, A., and Fadeel, B. (2014). Mitochondria released by cells undergoing TNF- α -induced necroptosis act as danger signals. *Cell Death Dis.* 5, e1312.
- Makki, K., Froguel, P., and Wolowczuk, I. (2013). Adipose tissue in obesity-related inflammation and insulin resistance: cells, cytokines, and chemokines. *ISRN Inflamm.* 2013, 139239.
- McLaughlin, T., Ackerman, S.E., Shen, L., and Engleman, E. (2017). Role of innate and adaptive immunity in obesity-associated metabolic disease. *J. Clin. Invest.* 127, 5–13.
- Mi, H., Huang, X., Muruganujan, A., Tang, H., Mills, C., Kang, D., and Thomas, P.D. (2017). PANTHER version 11: expanded annotation data from gene ontology and reactome pathways, and data analysis tool enhancements. *Nucleic Acids Res.* 45, D183–D189.
- Mooij, H.L., Bernelot Moens, S.J., Gordts, P.L., Stanford, K.I., Foley, E.M., van den Boogert, M.A., Wijtes, J.J., Hassing, H.C., Tanck, M.W., van de Sande, M.A., et al. (2015). Ext1 heterozygosity causes a modest effect on postprandial lipid clearance in humans. *J. Lipid Res.* 56, 665–673.
- Mousavi, S., Moradi, M., Khorshidahmad, T., and Motamedi, M. (2015). Anti-inflammatory effects of heparin and its derivatives: a systematic review. *Adv. Pharmacol. Sci.* 2015, 507151.
- Oduah, E.I., Linhardt, R.J., and Sharfstein, S.T. (2016). Heparin: past, present, and future. *Pharmaceuticals (Basel)* 9, 38.
- Ogden, C.L., Carroll, M.D., Lawman, H.G., Fryar, C.D., Kruszon-Moran, D., Kit, B.K., and Flegal, K.M. (2016). Trends in obesity prevalence among children and adolescents in the United States, 1988–1994 through 2013–2014. *JAMA* 315, 2292–2299.
- Orchard, R.C., Wilen, C.B., Doenck, J.G., Baldrige, M.T., McCune, B.T., Lee, Y.C., Lee, S., Pruett-Miller, S.M., Nelson, C.A., Fremont, D.H., and Virgin, H.W. (2016). Discovery of a proteinaceous cellular receptor for a norovirus. *Science* 353, 933–936.
- Orvedahl, A., McAllister, M.R., Sansone, A., Dunlap, B.F., Desai, C., Wang, Y.T., Balce, D.R., Luke, C.J., Lee, S., Orchard, R.C., et al. (2019). Autophagy genes in myeloid cells counteract IFN γ -induced TNF-mediated cell death and fatal TNF-induced shock. *Proc. Natl. Acad. Sci. USA* 116, 16497–16506.
- Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. (2017). Salmon provides fast and bias-aware quantification of transcript expression. *Nat. Methods* 14, 417–419.
- Patterson, R.E., Frank, L.L., Kristal, A.R., and White, E. (2004). A comprehensive examination of health conditions associated with obesity in older adults. *Am. J. Prev. Med.* 27, 385–390.
- Pham, A.H., McCaffery, J.M., and Chan, D.C. (2012). Mouse lines with photo-activatable mitochondria to study mitochondrial dynamics. *Genesis* 50, 833–843.
- Pirzgalska, R.M., Seixas, E., Seidman, J.S., Link, V.M., Sánchez, N.M., Mahú, I., Mendes, R., Gres, V., Kubasova, N., Morris, I., et al. (2017). Sympathetic neuron-associated macrophages contribute to obesity by importing and metabolizing norepinephrine. *Nat. Med.* 23, 1309–1318.
- Pollara, J., Edwards, R.W., Lin, L., Bendersky, V.A., and Brennan, T.V. (2018). Circulating mitochondria in deceased organ donors are associated with immune activation and early allograft dysfunction. *JCI Insight* 3, e121622.
- Qiu, Y., Nguyen, K.D., Odegaard, J.I., Cui, X., Tian, X., Locksley, R.M., Palmiter, R.D., and Chawla, A. (2014). Eosinophils and type 2 cytokine signaling in macrophages orchestrate development of functional beige fat. *Cell* 157, 1292–1308.
- Rao, R.R., Long, J.Z., White, J.P., Svensson, K.J., Lou, J., Lokurkar, I., Jedrychowski, M.P., Ruas, J.L., Wrann, C.D., Lo, J.C., et al. (2014). Meteorin-like is a hormone that regulates immune-adipose interactions to increase beige fat thermogenesis. *Cell* 157, 1279–1291.
- Raouf, R., van der Vlist, M., Willemen, H.L.D.M., Prado, J., Versteeg, S., Vos, M., Lockhorst, R., Pasterkamp, J., Khoury-Hanold, W., Meyaard, L., and Eijkelkamp, N. (2020). Macrophages transfer mitochondria to sensory neurons to resolve inflammatory pain. *bioRxiv*. <https://doi.org/10.1101/2020.02.12.940445>.
- Rebbbeck, C.A., Leroy, A.M., and Burt, A. (2011). Mitochondrial capture by a transmissible cancer. *Science* 331, 303.
- Reilly, S.M., and Saltiel, A.R. (2017). Adapting to obesity with adipose tissue inflammation. *Nat. Rev. Endocrinol.* 13, 633–643.
- Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* 43, e47.
- Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26, 139–140.
- Rutstein, D.D., Castelli, W.P., and Nickerson, R.J. (1969). Heparin and human lipid metabolism. *Lancet* 1, 1003–1008.
- Sasisekharan, R., and Venkataraman, G. (2000). Heparin and heparan sulfate: biosynthesis, structure and function. *Curr. Opin. Chem. Biol.* 4, 626–631.
- Scozzi, D., Ibrahim, M., Liao, F., Lin, X., Hsiao, H.M., Hachem, R., Tague, L.K., Ricci, A., Kulkarni, H.S., Huang, H.J., et al. (2019). Mitochondrial damage-associated molecular patterns released by lung transplants are associated with primary graft dysfunction. *Am. J. Transplant.* 19, 1464–1477.
- Singer, K., and Lumeng, C.N. (2017). The initiation of metabolic inflammation in childhood obesity. *J. Clin. Invest.* 127, 65–73.
- Sinha, P., Islam, M.N., Bhattacharya, S., and Bhattacharya, J. (2016). Inter-cellular mitochondrial transfer: bioenergetic crosstalk between cells. *Curr. Opin. Genet. Dev.* 38, 97–101.
- Sladek, R., Rocheleau, G., Rung, J., Dina, C., Shen, L., Serre, D., Boutin, P., Vincent, D., Belisle, A., Hadjadj, S., et al. (2007). A genome-wide association study identifies novel risk loci for type 2 diabetes. *Nature* 445, 881–885.
- Spees, J.L., Olson, S.D., Whitney, M.J., and Prockop, D.J. (2006). Mitochondrial transfer between cells can rescue aerobic respiration. *Proc. Natl. Acad. Sci. USA* 103, 1283–1288.
- Sun, K., Kusminski, C.M., and Scherer, P.E. (2011). Adipose tissue remodeling and obesity. *J. Clin. Invest.* 121, 2094–2101.
- The Gene Ontology Consortium (2017). Expansion of the Gene Ontology knowledgebase and resource. *Nucleic Acids Res.* 45, D331–D338.
- Torralba, D., Baixauli, F., and Sánchez-Madrid, F. (2016). Mitochondria know no boundaries: mechanisms and functions of intercellular mitochondrial transfer. *Front. Cell Dev. Biol.* 4, 107.

- Wang, L., Wang, S., and Li, W. (2012). RSeQC: quality control of RNA-seq experiments. *Bioinformatics* 28, 2184–2185.
- Wellen, K.E., and Hotamisligil, G.S. (2003). Obesity-induced inflammatory changes in adipose tissue. *J. Clin. Invest.* 112, 1785–1788.
- Wu, D., Molofsky, A.B., Liang, H.E., Ricardo-Gonzalez, R.R., Jouihan, H.A., Bando, J.K., Chawla, A., and Locksley, R.M. (2011). Eosinophils sustain adipose alternatively activated macrophages associated with glucose homeostasis. *Science* 332, 243–247.
- Wynn, T.A., Chawla, A., and Pollard, J.W. (2013). Macrophage biology in development, homeostasis and disease. *Nature* 496, 445–455.
- Zhao, S., Guo, Y., Sheng, Q., and Shyr, Y. (2014). Advanced heat map and clustering analysis using Heatmap3. *BioMed Res. Int.* 2014, 986048.
- Zhu, L., Zhang, J., Zhou, J., Lu, Y., Huang, S., Xiao, R., Yu, X., Zeng, X., Liu, B., Liu, F., et al. (2016). Mitochondrial transplantation attenuates hypoxic pulmonary hypertension. *Oncotarget* 7, 48925–48940.
- Zhu, C., Xu, P., He, Y., Yuan, Y., Wang, T., Cai, X., Yu, L., Yang, L., Wu, J., Wang, L., et al. (2017). Heparin increases food intake through AgRP neurons. *Cell Rep.* 20, 2455–2467.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-F4/80 MicroBeads UltraPure, mouse	Miltenyi Biotec	Cat#130-110-443
Anti-HS IgM (clone F58-10E4)	Amsbio	Cat#370255-1
Anti-mouse IgM-PE/Cy7 (clone RMM-1)	BioLegend	Cat#406514
Mouse anti-mouse CD64-PE-Dazzle594 (clone X54-5/7.1)	BioLegend	Cat#139304
Rat anti-mouse CD16/32 (clone 2.4G2)	BD Biosciences	Cat#553142
Rat anti-mouse CD206-PE/Cy7 (clone C068C2)	BioLegend	Cat#141720
Rat anti-mouse CD4-APC/Fire750 (clone RM4-5)	BioLegend	Cat#100568
Rat anti-mouse CD45-BUV395 (clone 30-F11)	BD Horizon	Cat#564279
Rat anti-mouse CD45.1-APC (clone A20)	BioLegend	Cat#110714
Rat anti-mouse CD45.2-BUV395 (clone 104)	BD Horizon	Cat#564616
Rat anti-mouse F4/80-APC (clone BM8)	BioLegend	Cat#123116
Rat anti-mouse iNOS-PE/Cy7 (clone CXNFT)	ThermoFisher/eBioscience	Cat#25-5920-82
Rat anti-mouse Ly6C-BV510 (clone HK1.4)	BioLegend	Cat#128033
Rat anti-mouse Ly6G-BV785 (clone 1A8)	BioLegend	Cat#127645
Rat anti-mouse SiglecF-BV421 (clone E50-2440)	BD Pharmingen	Cat#562681
Rat anti-mouse SiglecF-PE (clone E50-2440)	BD Pharmingen	Cat#552126
Rat anti-mouse/human ARG1-APC (clone A1exF5)	ThermoFisher/eBioscience	Cat#17-3697-82
Rat anti-mouse/human CD11b-BV650 (clone M1/70)	BioLegend	Cat#101259
Rat IgG2a kappa-APC (clone BR2a)	ThermoFisher/eBioscience	Cat#17-4321-81
Rat IgG2a kappa-PE/Cy7 (clone eBR2a)	ThermoFisher/eBioscience	Cat#25-4321-82
Rat-anti-mouse F4/80-PE (clone BM8)	BioLegend	Cat#123110
Bacterial and Virus Strains		
pXPR_011	Addgene	#59702
Mouse CRISPR Knockout Pooled Library (Brie) lentivirus	Addgene	#73633
lentiCRISPR v2	Addgene	#52961
Chemicals, Peptides, and Recombinant Proteins		
60% high fat diet, irradiated	Research Diets	Cat#D12492i
ACK Red Blood Cell Lysis Buffer	Gibco	Cat#A10492-01
Agencourt AMPure XP SPRI beads	Beckman Coulter	Cat#A63881
Bovine Serum Albumin	Pierce	Cat#23209
Brilliant Stain Buffer	BD Horizon	Cat#566349
CMTMR	ThermoFisher	Cat#C2927
Collagenase type II	Sigma-Aldrich	Cat#C6885
Contour handheld glucometer	Bayer	Cat#9545C
Coomassie Plus Reagent	Pierce	Cat#23236
Delbuco's Modified Eagle Media (DMEM) with 4.5 g/L D-glucose, L-glutamine, and no sodium pyrophosphate	Gibco	Cat#11965-084
DPBS	Gibco	Cat#14190-136
EDTA, 0.5M, pH 8.0	Invitrogen	Cat#AM9260G
ExTaq DNA polymerase	Clontech / Takara	Cat#RR001C
Fetal bovine serum (FBS)	Corning	Cat#35-010-CV
Fluoresbrite Polychromatic Red Microspheres, 1 micron	Polysciences	Cat#18660-5
Glucose	Sigma-Aldrich	Cat#G7528

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Grade I-A unfractionated heparin sodium salt, BioReagent	Sigma-Aldrich	Cat#H3149
Heparan sulfate proteoglycan	Sigma-Aldrich	Cat#H4777
Heparanase I	New England Biolabs	Cat#P0735S
Heparanase II	New England Biolabs	Cat#P0736L
Heparanase III	New England Biolabs	Cat#P0737L
Humulin R	Lilly	Cat#0002-8215-17
L-glutamine	Corning	Cat#25-005-CI
Lipopolysaccharide from E. coli O111:B4, BioXtra	Sigma-Aldrich	Cat#L4391-1MG
MitoID-Red	Enzo Life Sciences	Cat#ENZ-51007-500
MitoSOX Red	ThermoFisher	Cat#M36008
MitoTracker Red CMXRos	ThermoFisher	Cat#M7512
Penicillin-Streptomycin	Corning	Cat#30-002-CI
Puromycin	Gibco	Cat#A11138-03
Recombinant murine I Φ N- γ	R&D Systems	Cat#485-MI-100
Recombinant murine IL-4	R&D Systems	Cat#404-ML-100
Trypan blue	Sigma-Aldrich	Cat#T8154
Trypsin-EDTA	Gibco	Cat#15400-054
Ultracomp eBeads Compensation Beads	Invitrogen	Cat#01-2222-42
ZombieUV Fixable Viability Dye	BioLegend	Cat#423108
Critical Commercial Assays		
Cytofix/Cytoperm	BD Biosciences	Cat#554714
Mitochondria Isolation Kit, Mouse Tissue	Miltenyi Biotec	Cat#130-096-946
RNeasy Micro kit	Qiagen	Cat#74004
RNeasy Mini kit	Qiagen	Cat#74106
QIAmp DNA Mini Kit	Qiagen	Cat#51306
Deposited Data		
RNAseq dataset comparing mtD2- and mtD2+ macrophages from eWAT of MitoFat mice	GEO Database	GEO: GSE157462
Experimental Models: Cell Lines		
BV2 cells	Orchard et al., 2016	N/A
Experimental Models: Organisms/Strains		
<i>Adipoq</i> ^{Cre/+}	Jackson Labs	Stock no. 28020
C57BL/6J	Jackson Labs	Stock no. 664
CD45.1 (<i>Ptprc</i> ^{fl})	Jackson Labs	Stock no. 2014
<i>Ext1</i> ^{Flox/Flox}	Jackson Labs	Stock no. 9326
<i>Ext1</i> ^{Flox/Flox} ; <i>Lyz2</i> ^{Cre/+}	Generated in house	N/A
<i>Fpr1</i> ^{-/-}	Jackson Labs	Stock no. 32933
<i>Fpr1</i> ^{-/-} MitoFat	Generated in house	N/A
<i>Lyz2</i> ^{Cre/+}	Jackson Labs	Stock no. 4781
MitoFat (<i>mtD2</i> ^{Flox/+} ; <i>Adipoq</i> ^{Cre/+})	Generated in house	N/A
<i>mtDendra2</i> (i.e., PhAM, mtD2)	Jackson Labs	Stock no. 18397
<i>mtDendra2</i> ^{Flox/+} (i.e., PhAM Flox, mtD2 ^{Flox/+})	Jackson Labs	Stock no. 18385
Oligonucleotides		
sgRNA targeting <i>B3gal</i> ⁶ : CCGCGCTAAGGCCTTCCTGG	Integrated DNA Technologies	N/A
sgRNA targeting <i>B3gal</i> ³ : CTGGTCTCCTCTTTACACAC	Integrated DNA Technologies	N/A
sgRNA targeting <i>B4gal</i> ⁷ : CATCTATGTGCTCAACCAGG	Integrated DNA Technologies	N/A
sgRNA targeting <i>Ext1</i> : CATGGAGTCCTGCTTCGATT	Integrated DNA Technologies	N/A
sgRNA targeting <i>Ext2</i> : CCCTGAGTACAGAGAGGAAC	Integrated DNA Technologies	N/A
sgRNA targeting <i>Ext3</i> : CGCGGCTCTTCGAGGCCCTG	Integrated DNA Technologies	N/A

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
sgRNA targeting <i>Gfp</i> GAAGTTCGAGGGCGACACCC	Integrated DNA Technologies	N/A
sgRNA targeting <i>Hs6st1</i> CGCCGGTCTTCTGGATGTGC	Integrated DNA Technologies	N/A
sgRNA targeting <i>Papss1</i> TGCTACACTTTGGATGGTGA	Integrated DNA Technologies	N/A
sgRNA targeting <i>Slc35b2</i> CGGGTCTCCAGGTAAGAATA	Integrated DNA Technologies	N/A
sgRNA targeting <i>Ugdh</i> GCATTGTGCAGAACTCAAAT	Integrated DNA Technologies	N/A
sgRNA targeting <i>Ugp2</i> TCCAGGGCATGGAGATATCT	Integrated DNA Technologies	N/A
sgRNA targeting <i>Uxs1</i> : TCCACTCCGAGGTATATGG	Integrated DNA Technologies	N/A
TaqMan Gene Expression Assay Beta-actin FAM	Applied Biosystems	Mm00607939_s1
TaqMan Gene Expression Assay Ext1 FAM	Applied Biosystems	Mm00468769_m1
TaqMan Gene Expression Assay Ext2 FAM	Applied Biosystems	Mm00468775_m1
TaqMan Gene Expression Assay Hs6st1 FAM	Applied Biosystems	Mm01229698_s1
Software and Algorithms		
Prism 7	GraphPad Software	N/A
OxyMax	Columbus Instruments	N/A
STARS software	http://www.broadinstitute.org/rnai/public/software/index ; (Doench et al., 2016)	N/A
STAR version 2.0.4b	(Dobin et al., 2013)	N/A
Subread:featureCount v1.4.5	(Liao et al., 2014)	N/A
Sailfish version 0.6.13	(Patro et al., 2017)	N/A
RSeQC version 2.3	(Wang et al., 2012)	N/A
R/Bioconductor package EdgeR	(Robinson et al., 2010)	N/A
Limma	(Ritchie et al., 2015)	N/A
voomWithQualityWeights	(Liu et al., 2015)	N/A
GAGE	(Luo et al., 2009)	N/A
heatmap3	(Zhao et al., 2014)	N/A
Pathview	(Luo and Brouwer, 2013)	N/A
GO Consortium enrichment analysis tool	(Ashburner et al., 2000; Mi et al., 2017; The Gene Ontology, 2017)	N/A
Imaris software	BITPLANE	N/A
FlowJo v10	FlowJo, LLC	N/A

RESOURCE AVAILABILITY

Lead Contact

Requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Steven L. Teitelbaum (teitelbs@wustl.edu).

Materials Availability

All reagents are available from the Lead Contact under a material transfer agreement with Washington University in St. Louis.

Data and Code Availability

All data and code to understand and assess the conclusion of this research are available in the main text, supplementary materials, or GEO Database (accession number GSE157462).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mouse Models

C57BL/6J (stock number 664), CD45.1 (*Ptprc*^a; stock number 2014), *mtDendra2* (PhAM, referred to here as mtD2; stock number 18397), *mtDendra2*^{Flox/+} (PhAM Flox; referred to here as *mtD2*^{F/+} stock number 18385), *Adipoq*^{Cre/+} (stock number 28020), *Fpr1*^{-/-} (stock number 32933), *Ext1*^{Flox/+} (stock no 9326), and *Lyz2*^{Cre/+} (stock no 4781) mice were obtained from Jackson Laboratories (Bar Harbor, Maine). All mice were maintained on a C57BL/6J background, except *Ext1*^{Flox/Flox} and *Ext1*^{Flox/Flox}; *Lyz2*^{Cre/+} which were maintained on a mixed C57BL/6J;129 background and used as littermates. *mtD2*^{Flox/+} mice express mtD2 under control

of a ROSA26 Flox-STOP-Flox system, enabling expression of mtD2 in cells that express Cre recombinase. *mtD2^{Flox/Flox}* mice were crossed with *Adipoq^{Cre/+}* mice to generate *mtDendra2^{Flox/+} Adipoq^{Cre/+}* (MitoFat) and *mtDendra2^{Flox/+}* (control) mice. MitoFat mice were also crossed onto an *Fpr1^{-/-}* background. mtD2 mice were crossed to CD45.1 mice to generate CD45.1 mtD2 mice. Mice were genotyped using primers and protocols obtained from Jackson Laboratories or by Transnetyx. Mice were male or female, as indicated, and were age 6-26-weeks-old. All mice had *ad libitum* access to food water and were maintained in a specific-pathogen free facility with a 12h:12h light:dark cycle. Animals were randomly assigned to n=3-8 mice/group per experiment, except where indicated otherwise, and data represent at least 2 independent experiments throughout. Mice were euthanized for tissue harvest using carbon dioxide asphyxiation. All experiments were carried out under the guidelines of the Institutional Animal Care and Use Committee (IACUC) at Washington University in St. Louis and were performed under IACUC-approved protocols.

Cell Culture

BV2 microglia-derived cells (Orchard et al., 2016) were cultured in Delbuco's Modified Eagle Media (DMEM) with 4.5 g/L D-glucose, L-glutamine, and no sodium pyrophosphate (Gibco) that was additionally supplemented with 10% heat-inactivated fetal bovine serum (FBS), 2 mM L-glutamine, 100 U/mL Penicillin, and 100 U/mL Streptomycin (BV2 Media) and maintained at 37 °C with 5% CO₂. BV2 cells (1.0-2.0 × 10⁶) were seeded in 150 mm tissue culture-treated polystyrene plates, grown to ~80% confluence over 2-3 days before use or additional passaging.

METHOD DETAILS

Bone Marrow Chimeras

Male, 8-week-old mtD2 mice and C57BL/6J mice were lethally irradiated with 1,000 rad and one day later transplanted with 5.0 × 10⁶ bone marrow cells isolated from CD45.1 wildtype mice by intravenous administration. Bone marrow cells were isolated from the tibias and femurs and pooled from 5 mice. Recipient mice were allowed to engraft for 12 weeks before euthanization and tissue harvest.

Metabolic Cage Analyses

Ext1^{Flox/Flox};Lyz2^{Cre/+} mice and *Ext1^{Flox/Flox}* littermate control males (age 22-26 weeks old) were weighed and placed single-housed in a 16-metabolic cage Comprehensive Laboratory Animal Monitoring System (CLAMS) metabolic cages (Columbus Instruments, Columbus, OH). Mice were acclimated for approximately 16 hours followed by measurement for 24 hours from 07:00 (Zeitgeber time 0hr) to 06:59 the next day (Zeitgeber time 24hr). Mice were housed at room temperature on a 12h:12h light:dark cycle and arrayed in a staggered fashion to equally distribute the two groups from top to bottom and left to right. Mice had *ad libitum* access to food and water (hanging feeders and water bottles on load cells). Cumulative food intake over the 24 hr measurement period was determined. Activity was monitored using infrared laser/detector arrays on the X- and Y-axes. Oxygen consumption and carbon dioxide production were measured by indirect calorimetry using a zirconia O₂ sensor and CO₂ sensor at air flow rates of 0.90 L/min (18 sec line bleed followed by 2 sec measurement for each cage). Cages were measured individually in series (~5.5 min for each complete interval), with room air sampled between each interval to calculate O₂ consumption and CO₂ production. Heat was normalized to body weight. Data from 2 independent experiments were pooled for analysis with n=16 *Ext1^{Flox/Flox};Lyz2^{Cre/+}* mice and n=15 *Ext1^{Flox/Flox}* littermate controls from 7 litters.

Heparin Treatment

PBS or Grade I-A unfractionated heparin sodium salt (5 mg/kg body weight, equivalent to 900 U/kg; Sigma-Aldrich) in PBS were administered by intraperitoneal injection to MitoFat or *mtD2^{Flox/+}* control mice daily for 7 days at approximately 5:00 P.M. On the morning of day 7, an additional bolus of PBS or heparin was administered 30 min prior to tissue harvest for flow cytometric analysis.

Glucose Homeostasis Analyses

All blood glucose concentrations were measured using a Contour (Bayer) handheld glucometer with small droplets of blood obtained from the tail vein. For glucose tolerance tests (GTT), mice were fasted for 16 hours and then fasting blood glucose concentrations were determined and assigned as time point 0 min. A bolus of glucose (1 g glucose/kg body weight) in 0.45% saline was administered by intraperitoneal injection, and blood glucose concentrations were measured in duplicate or triplicate 20, 40, 60, 90, and 120 min later. For insulin tolerance tests (ITT), mice were fasted for 4 hours and then fasting blood glucose concentrations were determined and assigned as time point 0 min. A bolus of human insulin (Humulin R, 0.5 U/kg body weight) in 0.9% saline was administered by intraperitoneal injection, and blood glucose concentrations were measured in duplicate or triplicate 20, 40, and 60 min later.

Body Composition Analysis

Mice were weighed, and body composition was measured using an EchoMRI-100H 2n1 with a horizontal probe configuration (EchoMRI, Houston, TX).

Isolation of Immune Cells from Mouse Tissues

Epididymal white adipose tissue (eWAT), ovarian (o)WAT, inguinal (i)WAT, and/or brown adipose tissue (BAT) were finely minced and digested with 0.1% collagenase type II (Sigma-Aldrich, C6885) in DMEM at 37 °C for 60 min in an orbital shaker with rotation at 200 rpm while tilted at a 45° angle. Single cell suspensions of eWAT, iWAT, or BAT were passed through a 100 μ m nylon mesh filter, washed with 10 mL of Wash Media (DMEM with 5% FBS, 2 mM L-glutamine, and 100U/mL Penicillin-Streptomycin), and centrifuged at 500 \times g for 5 min at 4 °C. The floating adipocytes were aspirated and the stromal vascular fraction (SVF) pellet was resuspended in 1 mL ACK Red Blood Cell (RBC) Lysis Buffer (Gibco). After 3–5 min at room temperature, the RBC lysis reaction was quenched with 10 mL Wash Media, the cells were pelleted as above, and the pellet was resuspended in Wash Media for subsequent staining for flow cytometric analyses.

Flow Cytometry and Cell-Sorting

Cells were washed in 200 μ L PBS and then stained in 50 μ L of ZombieUV (1:600, BioLegend) in PBS and incubated on ice for 10 min covered in foil. The reaction was quenched with 200 μ L of FACS Buffer (PBS supplemented with 2.5% heat-inactivated PBS and 2 mM EDTA), and cells were pelleted as above. Cells were resuspended in 50 μ L of 10 μ g/mL FcBlock (rat anti-mouse CD16/32, clone 2.4G2, BD Biosciences) for 10 min covered in foil before addition of 50 μ L 2X stain cocktail (final concentration 1X, see below for antibody information and final dilutions) comprised of various fluorophore-conjugated antibodies in Brilliant Stain Buffer (BD Biosciences) containing 10 μ g/mL FcBlock. Cells were stained for 20–30 min on ice covered in foil and then washed 3 times in 200 μ L FACS Buffer. After the final resuspension, 25 μ L CountBright Absolute Counting Beads (ThermoFisher) were added for enumeration of cell numbers. For intracellular cytokine staining of BV2 cells with anti-ARG1-APC and anti-iNOS-PE/Cy7 (details below), cells were incubated with FcBlock as above and then fixed in 100 μ L of CytoFix/CytoPerm Buffer (BD Biosciences) for 30 min on ice while covered in foil. Cells were washed 3 times in 200 μ L of CytoPerm Wash (1X) per manufacturer's instructions and then stained in 1X CytoPerm Wash containing fluorophore-conjugated antibodies and 10 μ g/mL FcBlock for 1 hour on ice while covered in foil. Cells were washed 3 times in 200 μ L of CytoPerm Wash (1X), washed once in 200 μ L of FACS Buffer, and then resuspended in FACS Buffer for flow cytometry.

The following antibodies were used: rat anti-mouse SiglecF-PE (BD Pharmingen; clone E50-2440; 1:400), rat-anti-mouse F4/80-PE (BioLegend; clone BM8; 1:300), mouse anti-mouse CD64-PE-Dazzle594 (BioLegend; clone X54-5/7.1; 1:300), rat anti-mouse CD206-PE/Cy7 (BioLegend; clone C068C2; 1:200), rat anti-mouse iNOS-PE/Cy7 (eBioscience; clone CXNFT; 1:200), rat IgG2a kappa-PE/Cy7 (eBioscience; clone eBR2a; 1:200), rat anti-mouse CD45.1-APC (BioLegend; clone A20; 1:200), rat anti-mouse F4/80-APC (BioLegend; clone BM8; 1:300), rat anti-mouse/human ARG1-APC (eBioscience; clone A1exF5; 1:200), rat IgG2a kappa-APC (eBioscience; clone BR2a; 1:200), rat anti-mouse CD4-APC/Fire750 (BioLegend; clone RM4-5; 1:300), rat anti-mouse CD45-BUV395 (BD Horizon; clone 30-F11; 1:200), rat anti-mouse CD45.2-BUV395 (BD Horizon; Clone 104; 1:200), rat anti-mouse SiglecF-BV421 (BD Pharmingen; clone E50-2440; 1:400), rat anti-mouse Ly6C-BV510 (BioLegend; clone HK1.4; 1:400), rat anti-mouse Ly6G-BV785 (BioLegend; clone 1A8; 1:300), and rat anti-mouse/human CD11b-BV650 (BioLegend; clone M1/70; 1:400). For heparan sulfate (HS) quantification on cells, cells were stained with or without mouse anti-HS IgM antibody (amsbio; clone F58-10E4; 1:100) followed by a rat anti-mouse IgM-PE/Cy7 secondary antibody (BioLegend; clone RMM-1; 1:200).

All flow cytometry was performed on a 5-laser BD X20 or flow cytometer. Flow cytometric analyses were performed with FlowJo (version 10, Ashland, OR), and cells were gated on singlets and live cells. Macrophages were defined as CD45⁺ SiglecF⁺ CD11b⁺ F4/80⁺ CD64⁺ cells. Monocytes were defined as CD45⁺ SiglecF⁺ F4/80⁺ Ly6C^{hi} cells. Eosinophils were defined as SiglecF⁺ SSC^{hi} cells. Neutrophils were defined as CD11b⁺ Ly6G⁺ cells. Cell sorting of eWAT macrophages was performed on a BD Aria, and cells were sorted directly into RLT Buffer and then frozen on dry ice and stored at -80 °C until RNA extraction using the RNeasy Micro Kit (Qiagen).

Staining eWAT Macrophages with Mitochondria Dyes

eWAT SVF cells were isolated from male WT or MitoFat mice age 10–12 weeks old, as described above. Cells from 2 eWAT pads per mouse were resuspended in 400 μ L Wash Media and divided into 4 equal aliquots. To the aliquots were resuspended in 200 μ L Wash Media (unstained control) or 200 μ L Wash Media containing MitoD-Red (1:10,000 dilution; Enzo Life Sciences), CMX Red Rosamine (100 nM; ThermoFisher), or MitoSOX Red (5 μ M; ThermoFisher) per manufacturer protocols. Cells were incubated at 37°C with 5% CO₂ for 15 min and then washed with 200 μ L Wash Media twice and 200 μ L PBS before staining for flow cytometric analysis, as described above.

Adoptive Cell Transfer

eWAT SVF was obtained from n=3 C57BL/6J mice as described above and pooled for each independent experiment. RBC-depleted SVF cells were washed in 1 mL MACS Buffer (PBS containing 1% bovine serum albumin and 1 mM EDTA) and pelleted by centrifugation at 500 \times g for 5 min at 4 °C. The cells were resuspended in 90 μ L of MACS Buffer followed by addition of 10 μ L magnetic nano-beads conjugated to anti-F4/80 antibody (Miltenyi Biotec). After a 15-min incubation at 4 °C covered in foil, the SVF cells were passed through a 30 μ m nylon mesh and then an LS Column on a QuadroMACS magnet (Miltenyi Biotec). Cells were washed 3 times with 3 mL of MACS Buffer and then eluted in 5 mL MACS Buffer. Purity was confirmed by flow cytometry and was >95%. The cells were pelleted by centrifugation at 500 \times g for 5 min at 4 °C, and the cells were resuspended in 2 mL Wash Buffer containing

10 μ M CytoTracker Orange (also known as CMTMR; ThermoFisher; stock solution 10 mM in DMSO) and incubated at 37 °C for 30 min. The CMTMR-labelled cells were washed 2 times in 25 mL Wash Media and one time in 25 mL PBS. The washed cell pellet was resuspended in 300 μ L PBS for cell counting. Cells (approximately 5×10^4 per mouse) were administered by intraperitoneal injection into male or female mtD2 mice or male MitoFat mice and then imaged by intravital 2-photon microscopy 1–3 days after the adoptive transfer.

Intravital 2-Photon Microscopy

Mice were anesthetized by intraperitoneal injection of ketamine (50 mg/kg) and xylazine (10 mg/kg) and maintained with halved doses administered every hour. The mouse lower abdomen was carefully shaved with Nair (Church & Dwight Co.) and washed with PBS to remove the excess lotion. A small incision was made in the lower abdomen in order to expose the eWAT pad which was kept moist with PBS and carefully positioned under a coverslip that was screwed into place for imaging. During the acquisition, mouse status was closely monitored. Images were collected using a customized Leica SP8 2-photon microscope equipped with a 25X/0.95 NA water-dipping objective and a Mai Tai HP DeepSee Laser (Spectra-Physics) tuned to 880 nm. Fluorescence emission was separated by 3 high-efficiency dichroic mirrors cutting at 458, 495, and 560 nm (Semrock) and directly directed to 4 supersensitive external detectors. These detectors and a resonant galvo scanner with 12,000 Hz frequency very short dwell times under relatively low laser power. This prevents photo-conversion of the Dendra2 protein. 3D stacks consisting of between 21 and 31 planes (0.5 μ m step size) were captured every 30 seconds. Imaris software (BITPLANE) was used to reconstruct 3-dimensional (3-D) images, determine cellular localization with 3-D positional mapping, and generate movies derived from time-lapsed imaging.

RNA Sequencing and Analysis

RNA was extracted from sort-purified macrophages (see above) using Qiagen RNeasy Micro kit with on-column DNase I (Qiagen) digestion and then eluted in 14 μ L RNase/DNase-free water. Samples were prepared according to library kit manufacturer's protocol, indexed, pooled, and sequenced on an Illumina HiSeq. Basecalls and demultiplexing were performed with Illumina's bcl2fastq software and a custom python demultiplexing program with a maximum of one mismatch in the indexing read. RNA-seq reads were then aligned to the Ensembl release 76 top-level assembly with STAR version 2.0.4b (Dobin et al., 2013). Gene counts were derived from the number of uniquely aligned unambiguous reads by Subread:featureCount version 1.4.5 (Liao et al., 2014). Isoform expression of known Ensembl transcripts were estimated with Sailfish version 0.6.13 (Patro et al., 2017). Sequencing performance was assessed for the total number of aligned reads, total number of uniquely aligned reads, and features detected. The ribosomal fraction, known junction saturation, and read distribution over known gene models were quantified with RSeQC version 2.3 (Wang et al., 2012).

All gene counts were then imported into the R/Bioconductor package EdgeR (Robinson et al., 2010) and TMM normalization size factors were calculated to adjust for samples for differences in library size. Ribosomal genes and genes not expressed in the smallest group size minus one samples greater than one count-per-million were excluded from further analysis. The TMM size factors and the matrix of counts were then imported into the R/Bioconductor package Limma (Ritchie et al., 2015). Weighted likelihoods based on the observed mean-variance relationship of every gene and sample were then calculated for all samples with the voomWithQualityWeights (Liu et al., 2015). The performance of all genes was assessed with plots of the residual standard deviation of every gene to their average log-count with a robustly fitted trend line of the residuals. Differential expression analysis was then performed to analyze for differences between conditions and the results were filtered for only those genes with Benjamini-Hochberg false-discovery rate adjusted p-values less than or equal to 0.05.

For each contrast extracted with Limma, global perturbations in known Gene Ontology (GO) terms and KEGG pathways were detected using the R/Bioconductor package GAGE (Luo et al., 2009) to test for changes in expression of the reported log 2 fold-changes reported by Limma in each term versus the background log 2 fold-changes of all genes found outside the respective term. The R/Bioconductor package heatmap3 (Zhao et al., 2014) and Pathview (Luo and Brouwer, 2013) was used to display heatmaps or annotated KEGG graphs across groups of samples for each GO term or KEGG pathway (respectively) with a Benjamini-Hochberg false-discovery rate adjusted p-value less than or equal to 0.05. In addition, GO term representation among significantly down-regulated and significantly up-regulated genes were determined using GO Consortium enrichment analysis tool (Ashburner et al., 2000; Mi et al., 2017; The Gene Ontology, 2017).

Mitochondria and Bead Uptake Assays in BV2 Cells

BV2 cells were washed in phosphate buffered saline (PBS) and treated with 0.05% trypsin-EDTA (Gibco) in PBS for 10–30 min. Cells were washed in 1 volume of BV2 Media, pelleted by centrifugation at 500 $\times g$ for 5 min at 4 °C, resuspended in 10 mL BV2 Media, and cell density was counted using a hemocytometer with 0.1% trypan blue. Aliquots of 1.0×10^5 BV2 cells were used for co-culture with purified mitochondria (2 μ g, see below) or 1.0 μ m red latex beads (100 beads per cell, or MOI 100, Fluoresbrite Polychromatic Red Microspheres, Polysciences) in a volume of 200 μ L/well in polystyrene tissue-culture treated 96-well round-bottom plates for 90 min at 37 °C and 5% CO₂. Cells were washed three times in 200 μ L ice-cold BV2 Media and then twice in 200 μ L ice-cold FACS Buffer (PBS with 2.5% heat-inactivated FBS and 2.5 mM EDTA) before flow cytometric analyses. In some experiments, purified mtD2 mitochondria were incubated with 1 mg/mL heparan sulfate proteoglycan (HSPG, Sigma-Aldrich) for 30 min at room temperature prior to co-culture with BV2 cells. In other experiments, BV2 cells were incubated with heparanases I, II, and III (5 U/mL each, New England Biolabs) in DMEM for 1 hour at 37 °C and 5% CO₂ prior to co-culture with purified mtD2 mitochondria. In some experiments, BV2 cells

were stimulated with recombinant Interleukin (IL)-4 (20 ng/mL; R&D Systems) or recombinant interferon (IFN)- γ (10 ng/mL; R&D Systems) plus gamma-irradiated lipopolysaccharide (LPS, 1 ng/mL; Sigma-Aldrich) in BV2 Media for 24 hours prior to co-culture with 1 micron beads, purified mtD2 mitochondria, or being harvested in RLT Lysis Buffer for subsequent quantitative real-time polymerase chain reaction (PCR).

Quantitative Real-Time PCR

RNA was extracted from BV2 cells using the RNeasy Mini Kit (Qiagen) according to manufacturer instructions. Messenger (m)RNA was reverse transcribed using iScript Reverse Transcriptase Supermix (Biorad, cat no. 1708841). Expression of *Ext1*, *Ext2*, and *Hs6st1* was measured using TaqMan assays on a 7500 fast real-time PCR machine (Applied Biosystems).

Mitochondria Isolation

Mitochondria were isolated using the Mitochondria Isolation Kit, Mouse Tissue (Miltenyi Biotec). mtDendra2 mice were euthanized, the right atrium was punctured, and the mouse was perfused with 10 mL PBS (sterile, ice-cold) via the left ventricle. Perfused liver was harvested, finely minced, and homogenized in a 2 mL dounce homogenizer in 1 mL Lysis Buffer with 7-8 strokes. Tissue homogenate was transferred to a 15 mL conical tube containing 9 mL 1X Separation Buffer (SB) and mixed thoroughly. Anti-TOM40 antibody conjugated to magnetic nanobeads (50 μ L) was added and incubated in the dark at 4 $^{\circ}$ C with gentle rocking for 1 hour. The mixture was passed through a 30 μ m nylon mesh and then an LS Column associated with a QuadroMACS magnet (Miltenyi Biotec). The column was washed 3 times with 3 mL of 1X SB. Purified mitochondria were eluted with 1.5 mL of 1X SB and pelleted by centrifugation at 15,000 \times g for 2 min at 4 $^{\circ}$ C. The pellet was resuspended in 1 mL Storage Buffer (Miltenyi Biotec), pelleted again as above, resuspended in 1 mL Storage Buffer, and held on ice. Mitochondria protein concentration was determined using 10 μ L of purified mitochondria and 300 μ L Coomassie Plus Reagent (Pierce) alongside a standard curve of Bovine Serum Albumin (BSA, Pierce). Absorbance was measured at 450 nm using a BioRad iMark Microplate Reader.

Brie CRISPR Library Preparation

Lentiviral libraries were synthesized, cloned, and produced as previously described (Doench et al., 2016). Cas9 activity was confirmed in stably-expressing BV2 cells (BV2-Cas9) using pXPR_011 (Addgene #59702) EGFP expression on a FACS Calibur instrument (BD Biosciences) (Orchard et al., 2016; Orvedahl et al., 2019). The lentiviral library was titered by transducing 3×10^6 BV2 cells with variable volumes of lentivirus (0, 10, 20, 40, 80, or 160 μ L) in 2 mL total volume of cell culture media in 50 mL conical tubes. The cells were spinoculated for 2 hr at 1000g at 30 $^{\circ}$ C. After spinoculation, cells were resuspended in cell culture media and 5×10^5 cells were plated per 15 cm dish. Media was exchanged 48 hr later with D10 or puromycin (Puro)-containing media, and cells were counted 48-72 hours later. A standard curve was then interpolated to determine lentiviral concentration needed for transduction efficiency of 20-30%. For library production, 1.6×10^8 BV2-Cas9 cells at 1.5×10^6 cells/mL were spinoculated with Brie lentivirus (Addgene #73633) sufficient to transduce 25% of cells. This was sufficient for 500x coverage for each of the approximately 80,000 sgRNAs. Following spinoculation, cells were resuspended, and plated at a density of 5×10^5 cells in 25 mL D10 in 15 cm dishes. Forty-eight hours after transduction, media was exchanged with puromycin containing media and cells were cultured for an additional three days prior to pooling, aliquoting, and freezing in liquid nitrogen.

Genome-Wide CRISPR Knockout Screen, Sequencing, and Analysis

An aliquot of 4×10^7 BV2-Brie cells were seeded in 20 \times 150 mm tissue-culture treated dishes with 2×10^6 cells in 25 mL BV2 Media per dish and cultured for 48 hours, as described above. The cells were harvested as described above and counted using a hemocytometer. An aliquot of 1.5×10^8 cells were cultured in 150 mL of BV2 Media containing 6.0 mg (40 μ g/mL) of purified mitochondria from livers of $n=5$ mtD2 mice for 90 min at 37 $^{\circ}$ C with gentle swirling each 15 min. Cells were pelleted by centrifugation at 500 \times g for 5 min at 4 $^{\circ}$ C, washed in 50 mL of BV2 Media three times before washing once in 50 mL of PBS. Cells were stained with ZombieUV and rat anti-mouse F4/80-APC, as described above, in filter-sterilized FACS Buffer, and cells were sort-purified as described above. An aliquot of 1×10^7 cells was exposed to no mitochondria to determine a negative gate for sorting. The mitochondria-exposed pool of cells was held on ice, and mtD2-negative cells were sorted using an Aria I into sterile 2 mL sterile polypropylene tubes. Genomic DNA was isolated from the mtD2-negative BV2-Brie cells and an unsorted population 2×10^8 input cells using the QIAmp DNA Mini Kit according to manufacturer instructions.

Enrichment of sgRNA guides in mtD2-negative BV2 cells was determined by Illumina sequencing and STARS analysis, as previously described (Doench et al., 2016). Briefly, genomic DNA from mtD2-negative and input cells were aliquoted into multiple wells of 96-well plates with up to 10 μ g DNA in a maximum 50 μ L volume. Polymerase chain reaction (PCR) master mix (40 μ L) containing ExTaq DNA polymerase (Clontech), ExTaq buffer, dNTP, P5 stagger primer, and water was added along with 10 μ L of a barcoded primer, making a reaction volume of 100 μ L. Samples were amplified with the following program: 95 $^{\circ}$ C (1 min) followed by 28 cycles of 94 $^{\circ}$ C for 30 sec, 52.5 $^{\circ}$ C for 30 sec, 72 $^{\circ}$ C for 30 sec, with a final 10 min at 72 $^{\circ}$ C. PCR product was purified with Agencourt AMPure XP SPRI beads per the manufacturer's protocol (Beckman Coulter). Samples were sequenced on an Illumina HiSeq 2000. Barcodes in the P7 primer were de-convoluted and the sgRNA sequence was mapped to the reference of sgRNAs in the Brie library. To normalize for different numbers of reads per condition, read counts per sgRNA were normalized to 1×10^7 total reads per sample. This normalized value was then log-2 transformed. sgRNAs that were not sequenced were arbitrarily assigned a read count of 1.

sgRNA frequencies were analyzed using STARS software, which is available at <http://www.broadinstitute.org/rnai/public/software/index> (Doench et al., 2016). STARS computes a score for each gene of a rank-ordered sgRNA hits that was above 10% of total sequenced sgRNAs. A STAR score was only assigned to genes that scored above this threshold in at least two of the guides targeting that gene. In addition to the STAR score, we computed false discovery rates (FDR) to correct for multiple testing. FDR estimates the probability that the null hypothesis is true (i.e., a gene is not a hit). Thus, the genes with the lowest FDRs are the most likely to be true hits.

Generation of BV2 Knockout Cells

Knockout (KO) cell lines in BV2 cells with lentiCRISPR v2 (Addgene #52961) lentivirus containing Cas9, a puromycin resistance gene, and the following CRISPR sgRNA guide sequences: *Hs6st1* (CGCCGGTCTTCTGGATGTGC); *Uxs1* (TCCACTTCCGAGGTATATGG); *Slc35b2* (CGGGTCTCCAGGTAAGAATA); *Papss1* (TGCTACACTTTGGATGGTGA); *Ugp2* (TCCAGGGCATGGAGATATCT); *B3gat3* (CTGGTCTCCTCTTACACAC); *Extl3* (CGCGGCTCTTCGAGGCCCTG); *B4galt7* (CATCTATGTGCTCAACCAGG); *B3galt6* (CCGC GCTAAGGCCTTCCTGG); *Ext1* (CATGGAGTCCTGCTTCGATT); *Ext2* (CCCTGAGTACAGAGAGGAAC); and *Ugdh* (GCATTGTGCA GAACCAAAT). As a control, sgRNA directed against *GFP* (GAAGTTCGAGGGCGACACC) was used. Transduced cells were cultured at 37 °C and 5% CO₂ for 48 hours and then selected with puromycin (2.5 µg/mL) for 7 days. Clonal KO cell lines were generated by limiting dilution, and gene deletion was verified by deep sequencing. Clonal KO cells were propagated in BV2 Media, and aliquots of cells were frozen at -80 °C in heat-inactivated FBS containing 10% DMSO and stored at -80 °C till use. Thawed cells were plated in 150 mm dishes and cultured for 2 days to approximately 80% confluence, and aliquots of cells were co-cultured with purified mtD2 mitochondria or red latex beads, as described above. Flow cytometry was used to quantify the frequency of cells that were mtD2⁺ or bead⁺ using a BD X20 flow cytometer.

WAT Co-culture System

WAT from donor CD45.1 or CD45.2 mtD2 mice fed a normal chow or HFD were co-cultured with equal masses of WAT from recipient congenically disparate CD45.1 or CD45.2 WT mice fed a chow or HFD. Equal masses of donor and recipient WAT (within 0.01–0.03 g) were finely minced together in DMEM containing 0.1% collagenase II at 37 °C for 120 min with rotation at 200 rpm at 45° angle. The SVF was isolated and stained for flow cytometric analyses, as described above. The frequency of recipient macrophages that were mtD2⁺ was quantified.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical Analyses

Data are expressed as mean ± standard error of the mean (SEM). Statistical significance for normally distributed data was determined using Student's t-tests for comparisons of 2 groups or analysis of variance (ANOVA) followed by Fisher LSD post-hoc tests for comparisons of 3 or more groups. For metabolic cage analyses, ANOVA with repeated measures and Fisher LSD post-hoc tests were used. Significance was set at P<0.05. Statistical analyses were performed with Prism 7 (GraphPad Software), unless otherwise indicated.


Supplemental Information

**Intercellular Mitochondria Transfer
to Macrophages Regulates White Adipose
Tissue Homeostasis and Is Impaired in Obesity**

Jonathan R. Brestoff, Craig B. Wilen, John R. Moley, Yongjia Li, Wei Zou, Nicole P. Malvin, Marina N. Rowen, Brian T. Saunders, Hongming Ma, Madison R. Mack, Barry L. Hykes Jr., Dale R. Balce, Anthony Orvedahl, Jesse W. Williams, Nidhi Rohatgi, Xiaoyan Wang, Michael R. McAllaster, Scott A. Handley, Brian S. Kim, John G. Doench, Bernd H. Zinselmeyer, Michael S. Diamond, Herbert W. Virgin, Andrew E. Gelman, and Steven L. Teitelbaum


SUPPLEMENTAL FIGURES

Supplemental Fig. S1




Supplemental Figure S1. Bone marrow transplant chimerism and immune cell identification, related to Figure 1. (A) Chimerism and tissue-specific macrophage gating strategy in eWAT, iWAT, and BAT. Pre-gated on singlet, live cells. Data are expressed as mean ± SEM. **(B)** Gating strategy to identify various immune cell populations in eWAT.

Supplemental Fig. S2




Supplemental Figure S2. Heat map of top 20 differentially expressed genes in mtD2⁺ and mtD2⁻ macrophages that had or had not acquired mitochondria from adipocytes, related to Figure 3. (A) Singlet, live, CD45⁺ SiglecF⁻ CD64⁺ F4/80⁺ macrophages from eWAT of MitoFat mice were sort-purified on the basis of whether they were mtD2⁺ or mtD2⁻. The top 20 most differentially expressed genes in eWAT macrophages with adjusted P<0.05 are shown.

Supplemental Fig. S3




Supplemental Figure S3. FPR1 is not required for mitochondria transfer from adipocytes to macrophages, related to Figure 4. (A) eWAT SVF was isolated and singlet, live, CD45⁺ SiglecF⁻ cells were pre-gated. CD64⁺ F4/80⁺ macrophages were identified *Fpr1*^{+/+} MitoFat and *Fpr1*^{-/-} MitoFat mice and macrophage frequencies are shown. (B) Frequencies of mtD2⁺ macrophages in eWAT macrophages of *Fpr1*^{+/+} MitoFat and *Fpr1*^{-/-} MitoFat mice. N=7-8 mice/group, Student's t-tests performed. No significant differences were found. Data are expressed as mean ± SEM.

Supplemental Fig. S4



Supplemental Figure S4. Characterization of BV2 cell mitochondria uptake, related to Figure 4. **(A)** 10^5 BV2 cells suspended in 200 μ L culture media were cultured with various concentrations of liver-derived mtD2 mitochondria for 90 min. The frequencies of mtD2⁺ BV2 cells were measured for flow cytometry. Data are expressed as mean \pm SEM from n=3 independent experiments. **(B)** Representative flow cytometry plots used for sort-purifying mtD2-negative BV2 cells in the genome-wide CRISPR knockout screen. BV2-Brie cells were treated without (Control) or with 40 μ g/mL purified mtD2 mitochondria for 90 min and sort-purified. Control cells were used to define a negative gate. That gate was then applied to the mtD2-exposed cells for sorting of mtD2-negative cells. The frequencies of mtD2-negative cells were 3.75% in Replicate 1 and 3.50% in Replicate 2 of the screen. **(C)** Representative flow cytometry histograms of sulfate levels on BV2-Cas9 cells with sgRNA targeting green fluorescent protein (control) or Ext1.

Supplemental Fig. S5



Supplemental Figure S5. Polarization of BV2 cells towards an M1-like phenotype impairs mitochondria uptake while increasing phagocytosis of beads, related to Figure 5. BV2 cells (4×10^6) were treated with phosphate buffered saline (PBS), 20 ng/mL interleukin (IL)-4, or 10 ng/mL interferon (IFN)- γ plus 1 ng/mL lipopolysaccharide (LPS) for 24 hours. Suspensions of polarized BV2 cells (1×10^5) were then co-cultured with 2 μ g purified mtD2 mitochondria or 1×10^7 polychromatic red 1.0 μ m latex beads (multiplicity of infection, MOI 100) in a volume of a 200 μ L for 90 min at 37 $^{\circ}$ C. Intracellular cytokine stain for (A) Arginase 1 (ARG1) or (B) Inducible nitric oxide synthase (iNOS) proteins. Mean fluorescence intensities (MFI) were determined, and MFI for isotype controls was subtracted. Data are expressed as mean \pm SEM from n=3 independent experiments. One-way ANOVA with Fisher's LSD post-hoc test, ***P<0.001. (C) Representative flow cytometry plots of bead uptake. (D) Representative flow cytometry plots of mtD2 mitochondria uptake.